



# Why electrohypersensitivity and related symptoms are caused by non-ionizing man-made electromagnetic fields: An overview and medical assessment

Dominique Belpomme<sup>a,b,\*</sup>, Philippe Irigaray<sup>b</sup>

<sup>a</sup> Medical Oncology Department, Paris University, Paris, France

<sup>b</sup> European Cancer and Environment Research Institute (ECERI), Brussels, Belgium

## ARTICLE INFO

### Keywords:

EHS  
Electrohypersensitivity  
Electromagnetic field  
Extremely low frequency  
Microwaves  
Pathophysiological mechanism  
Radiofrequency

## ABSTRACT

Much of the controversy over the cause of electrohypersensitivity (EHS) lies in the absence of recognized clinical and biological criteria for a widely accepted diagnosis. However, there are presently sufficient data for EHS to be acknowledged as a distinctly well-defined and objectively characterized neurologic pathological disorder. Because we have shown that 1) EHS is frequently associated with multiple chemical sensitivity (MCS) in EHS patients, and 2) that both individualized disorders share a common pathophysiological mechanism for symptom occurrence; it appears that EHS and MCS can be identified as a unique neurologic syndrome, regardless their causal origin. In this overview we distinguish the etiology of EHS itself from the environmental causes that trigger pathophysiological changes and clinical symptoms after EHS has occurred. Contrary to present scientifically unfounded claims, we indubitably refute the hypothesis of a nocebo effect to explain the genesis of EHS and its presentation. We as well refute the erroneous concept that EHS could be reduced to a vague and unproven “functional impairment”. To the contrary, we show here there are objective pathophysiological changes and health effects induced by electromagnetic field (EMF) exposure in EHS patients and most of all in healthy subjects, meaning that excessive non-thermal anthropogenic EMFs are strongly noxious for health. In this overview and medical assessment we focus on the effects of extremely low frequencies, wireless communications radiofrequencies and microwaves EMF. We discuss how to better define and characterize EHS. Taken into consideration the WHO proposed causality criteria, we show that EHS is in fact causally associated with increased exposure to man-made EMF, and in some cases to marketed environmental chemicals. We therefore appeal to all governments and international health institutions, particularly the WHO, to urgently consider the growing EHS-associated pandemic plague, and to acknowledge EHS as a mainly new real EMF causally-related pathology.

## 1. Introduction

We have previously published evidence that a) electrohypersensitivity (EHS) is a distinct newly identified and objectively characterized neurologic pathological disorder which can be clinically diagnosed, and treated using peripheral blood and urine molecular

biomarkers and cerebral imaging (Belpomme and Irigaray, 2020); b) EHS and Multiple Chemical Sensitivity (MCS) are possibly associated in EHS patients, both presenting similar clinical presentation and biological and radiological abnormal changes, therefore EHS and MCS could in fact be two etiopathogenic disorders of a unique common pathological syndrome (Belpomme et al., 2015, 2016); c) EHS and MCs are both

**Abbreviations:** BBB, Blood brain barrier; CNS, Central nervous system; ECG, Electrocardiogram; EEG, electroencephalogram; EHS, Electrohypersensitivity; ELF, Extremely low-frequency; EMF, Electromagnetic field; EMG, Electromyogram; EMR, Electromagnetic Radiation; ESP, Electric skin potential; GSM, Global System for Mobile telecommunication; HRV, Heart rate variability; HSP, heat shock protein; IEI, Idiopathic environmental intolerance; IEI-EMF, Idiopathic environmental intolerance attributed to EMF; MCS, Multiple chemical sensitivity; MF, Magnetic field; MT, Mobile telephony; MW, Microwaves; OS, Oxidative stress; PET, Positron emission tomography; RBC, Red blood cells; RF, Radio frequency; SCBF, Skin capillary blood flow; VDT, Visual display terminal; WC, Wireless Communication; WHO, World Health Organization; WiFi, Wireless fidelity; WLAN, Wireless Local Area Network (for example WIFI).

\* Corresponding author. ARTAC, 57/59 rue de la Convention, 75015, Paris, France.

E-mail address: [contact.belpomme@gmail.com](mailto:contact.belpomme@gmail.com) (D. Belpomme).

<https://doi.org/10.1016/j.envres.2022.113374>

Received 13 December 2021; Received in revised form 30 March 2022; Accepted 25 April 2022

Available online 7 May 2022

0013-9351/© 2022 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

associated with detectable low grade inflammation (Belpomme et al., 2015) and oxidative stress (Irigaray et al., 2018a) with possible consequent blood brain barrier (BBB) opening (Belpomme and Irigaray, 2020) as in Alzheimer diseases (Heneka and O'Banion, 2007; Bell and Zlokovic, 2009; Erickson and Banks, 2013) and in other chronic pathological disorders (Patel and Frey, 2015) and d) EHS is associated with brain neurotransmitters abnormal concentrations (Belpomme and Irigaray, 2020) as in laboratory animals exposed to man-made electromagnetic fields (EMF) (Hu et al., 2021).

In a recent scientific international consensus report molecular biomarkers and imaging have been recognized to be of critical value to study EHS by many scientists (Belpomme et al., 2021). In addition, as emphasized in this report, a clear distinction has been made between the causal origin of EHS itself (its etiology) and the daily environmental causes that trigger pathophysiological changes and clinical symptoms in EHS patients after EHS has occurred (its pathogenesis). A pending question is however the role of EMF exposure, both in triggering clinical symptoms and biological changes, and in causing EHS itself. At present, the lack of clear answer to these two questions may explain why most mainstream medical, sanitary and societal bodies still believe that there is not sufficient scientific proof to assert that the clinical symptoms experienced by EHS self-reported patients are really caused by EMF exposure; nor that EHS genesis could be the consequence of excessive man-made EMF exposure. Additionally, since the World Health Organization (WHO) officially stated in 2005 (WHO, 2005) and more recently in 2014 (WHO, 2014), that EHS is a “disabling condition” associated with “non-specific symptoms that lack apparent toxicological or physiological basis or independent verification” and that there are “no clear diagnosis criteria”; it is widely accepted that EHS cannot be diagnosed medically and is not causally related to EMF exposure.

The uncertainty of provocation studies testing the existence of a positive correlative effect of EMF exposure versus sham exposure in EHS patients explain why the cause of symptomatic occurrence is still debated among scientists, some of them refuting the possibility of a causal effect of EMF in triggering symptoms not only in EHS patients (Levallois, 2002; Röösli, 2008; Röösli et al. 2010a, b) but also in healthy people (Baliatsas et al., 2015); some others postulating that EHS is of psychologic origin, i.e. a psychosomatic disease (Rubin et al., 2010, 2011); while still others contrary to the present WHO statements even question the existence of EHS itself (Leszczynski, 2021).

Recalling the historical main scientific research steps and the international institutional statements concerning EHS and MCS, we would like here to summarize how man-made EMF exposure and in some cases marketed environmental chemicals can really trigger symptoms in EHS patients, that exposure to non-thermal man-made EMF are objectively noxious for healthy people and that the etiology of EHS is in fact mainly causally related to man-made EMF exposure in genetically (or epigenetically) susceptible people.

There are indeed three scientific questions to address: a) what is the state of research on EHS pathogenesis b) how can we define hypersensitivity in EHS patients; and c) what is the etiology of EHS in genetically (or epigenetically) susceptible subjects and how it may be generated.

Before answering these questions we would like to emphasize that any causality determination must satisfy the following four WHO causality criteria: a) “the existence of biological effects and health hazards can only be established when research results are replicated in independent laboratories or supported by related studies”; b) “there is agreement with accepted scientific principles”; c) “the underlying mechanism is understood”; d) and finally “a dose-response can be established” (WHO, 2006).

Taking into account these four criteria we disclose and discuss here the present scientific state-of-the-art about the three above distinct scientific questions.

We would like as much as possible to attempt to distinguish the effect of extremely low electromagnetic frequency (ELF) (50–60 Hz), Wireless communication (WC) radiofrequency (RF) (3 kHz to 300 GHz) and WC

microwave (MW) EMF (300 MHz–300 GHz); which are presently used for different societal purposes. We would like also to specify that RF/MW electromagnetic radiation (EMR) used as carrier signals (300 kHz–300 GHz) is modulated by ELF EMR (3 Hz–3000 Hz) in order to transmit increasing amounts of information (Panagopoulos, 2019).

## 2. Historical scientific and institutional background

The term electromagnetic hypersensitivity which is commonly named electrohypersensitivity (EHS) was first proposed in 1991 by William Rea to identify the pathological condition of patients reporting health effects while being experimentally exposed to RF EMF versus sham and being compared to healthy controls in a controlled environment (Rea et al., 1991). This term was then re-used in 1997 in a report provided by a European group of scientific experts for the European Commission to clinically describe this unusual pathological condition, which posit EMF exposure as symptomatic trigger (Bergqvist and Vogel, 1997). In 2004, because of the seemingly worldwide prevalence increase in EHS, WHO organized an international scientific workshop in Prague to define and characterize EHS. Although not acknowledging EHS as being caused by EMF exposure, due to a lack of available correlation studies, the Prague working group clearly defined EHS as “a phenomenon where individuals experience adverse health effects while using or being in the vicinity of devices emanating electric, magnetic, or electromagnetic fields” (Mild et al., 2006). According to a previous 1996 WHO-sponsored International Program on Chemical Safety (IPCS)-related conference in Berlin on MCS (Report of the Workshop on Multiple Chemical Sensitivities, 1996), it was recommended to qualify such unknown new environmental pathological conditions under the term of “idiopathic environmental intolerance (IEI)”. Thus, following the 2004 Prague workshop, instead of using the term EHS, it was recommended to use the term idiopathic environmental intolerance attributed to EMF (IEI-EMF) to name this particular pathological condition, because of the lack of a proven causal link between EHS and EMF exposure, and no known pathophysiological mechanism linking EMF exposure with clinical symptoms. However, because the term EHS was in common use worldwide, WHO officially acknowledged also EHS as an adverse health condition in its 2005 fact sheet N°296 (WHO, 2005); and in its 2014 fact sheet N°193 which further reports on public health and mobile phone use, claiming again a lack of proven causal link between the emission of EMF from mobile phones and health effects, and that there is no proven underlying pathophysiological mechanism accounting for such effects (WHO, 2014). But it was already shown that mobile phones and more generally WC EMFs can cause clinical symptoms (NIEHS, 1998; Chia et al., 2000; Santini et al., 2002, 2003; and others), Oxidative Stress (OS) and DNA damage (Lai and Singh, 1995; Ivancsits et al., 2002, 2003; Diem et al., 2005; Panagopoulos et al., 2007; De Iulius et al., 2009; Phillips et al., 2009), while the biophysical mechanism of action was also already suspected (Panagopoulos et al., 2002).

Indeed since the 2005 and 2014 WHO official statements; much clinical, biological, and biophysical progress has been made to confirm previous data and to better understand the biophysical and biological processes of the noxious effects of EMFs (Panagopoulos et al., 2015a, 2021; Yakymenko et al., 2016; Lai 2019; 2021) and their pathophysiological significance on human health (Belpomme et al. 2015, 2018; Irigaray et al., 2018a); more particularly to identify and characterize EHS as a new pathological disorder (Belpomme and Irigaray, 2020). Such progress on EMF effects and EHS genesis was summarized in an international consensus meeting held in 2015 at the Royal Belgium Academy of Medicine in Brussels and published in a special issue of the journal *Reviews on Environmental Health* (Carpenter and Belpomme, 2015). Table 1 summarizes the historical scientific steps and WHO statements concerning MCS and EHS acknowledgment.

**Table 1**

The different historical steps to identify and qualify EHS and MCS, including WHO official statements, statements from WHO-sponsored meetings, and other scientific consensus meetings and reports.

|      |                                                                                                                                     |                                                                  |
|------|-------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------|
| 1962 | First identification and description of MCS                                                                                         | Randolph (1962)                                                  |
| 1991 | First identification and description of EHS                                                                                         | Rea et al. (1991)                                                |
| 1996 | Berlin WHO-sponsored workshop: MCS classified as idiopathic environmental intolerance (IEI)                                         | Report of the Workshop on Multiple Chemical Sensitivities (1996) |
| 1997 | Stockholm possible health implication of EMF exposure: a report prepared by a European group of experts for the European Commission | Bergqvist and Vogel (1997)                                       |
| 1999 | Atlanta (US), definition of MCS:1999 consensus meeting                                                                              | Bartha et al. (1999)                                             |
| 2004 | Prague WHO sponsored workshop: identification of idiopathic environmental intolerance attributed to EMF                             | Mild et al. (2006)                                               |
| 2005 | WHO fact sheet n° 292 aiming at defining EHS                                                                                        | WHO (2005)                                                       |
| 2014 | WHO fact sheet n° 193: EMF and Public Health; mobile phone                                                                          | WHO (2014)                                                       |
| 2015 | Brussels: Fourth Paris Appeal Colloquium; a focus on EMF and EHS                                                                    | Carpenter and Belpomme (2015)                                    |
| 2021 | The critical Importance of molecular biomarkers and imaging in the study of EHS. A scientific consensus international report        | Belpomme et al. (2021)                                           |

### 3. Symptomatic and biological triggers in EHS patients

Clinical symptoms presumably related to MW exposure were initially reported by Soviet scientists (Dodge, 1969; Carpenter, 2015). They consisted of headaches, fatigue, loss of appetite, insomnia, loss of concentration and short-term memory, transient cardiovascular dysfunction and labile emotional behavior. Some or all of these symptoms were described in particular in people exposed to microwave radar equipment. During the Soviet period, this symptomologic description was not acknowledged by western scientists. However in a 1972 revised document the US Naval Medical Research Institute was able to count more than 2500 references on the biological and clinical response to radio-frequency radiation (RFR) or microwave radiation published up to April 1972 in the world scientific literature (Glaser, 1972).

In 1979 the clinical symptoms reported to be caused by microwaves were recorded in the framework of a new clinical syndrome named the “microwave syndrome” (Pollack, 1979). This particular clinical syndrome considered to be caused by microwaves in exposed workers was described to involve the nervous system and to be characterized clinically by symptoms such as fatigue, headaches, dysesthesia and various autonomic dysfunctions. This microwave syndrome is symptomatically tantamount to the experimentally identified pathological disorder termed hypersensitivity to EMF (i.e. EHS) by William Rea in 1991 (Rea et al., 1991).

A first approach in describing the adverse health effects possibly associated with exposure to man-made EMFs was made in Sweden in 1984 by Ulf Bergqvist, who reported in a well-documented overview article the clinical symptoms occurring in people using Visual display terminal (VDT) (Bergqvist, 1984). Recorded symptoms included eye problems, ocular disturbance with change in visual performance, musculoskeletal discomfort, facial skin rashes, stress and psychological distress involving particularly mood disturbance, and adverse pregnancy outcomes. Although it was shown that there was an increased number and mobilization of mast cells in the skin of normal volunteers using VDT or television (TV) (Johansson et al., 2001), suggesting that these adverse health effects could be EMF-related; no clear causal relationship could be established between symptom occurrence and VDT- or TV-related EMF exposure. Thus, this observational study could not

relate specifically any symptom occurrence to EMF exposure.

Following this VDT study, Ulf Bergqvist and Evi Vogel, with other European scientific experts working for the European Commission conducted a multinational questionnaire-based survey and reported in 1997 that patients who claim to be EHS frequently have “neurasthenia” symptoms, headache and skin symptoms, and less frequently sleep disturbance and anxiety (Bergqvist and Vogel, 1997). However, again, these symptoms were considered non-specific and not causally related to EMF exposure. In fact, this large multinational questionnaire-based survey was unable to clinically define the real symptomatic picture presented by so-called EHS patients and its possible connection with EMF exposure.

However, in 1998, it was reported by the US National Institute of Environmental Health Sciences that health effects could be caused by exposure to powerline frequency (50–60 Hz) electric and magnetic fields (NIEHS, 1998), while in 2000 an increased prevalence of headache among mobile phone users was observed in Singapore (Chia et al., 2000). Then in 2002 Roger Santini in France described the clinical symptoms ascribed to mobile phone use in a French engineering school (Santini et al., 2002), and a year later those ascribed to Mobile Telephony (MT) based station proximity (Santini et al., 2003).

In fact, many studies focused on the symptomatic risk in ELF, RF or MW EMF-exposed people in the general population, but not specifically in EHS self-reported patients. All these general population-based studies were based on telephone survey or mailed or web-based questionnaires. Moreover, most of these studies in the general population investigated one or few self-reported symptoms such as headache (Chia et al., 2000; Milde-Busch et al., 2010; Sudan et al., 2012; Auvinen et al., 2019), tinnitus (Frei et al., 2012; Medeiros and Sanchez, 2016; Auvinen et al., 2019), sleep disturbance (Hutter et al., 2006; Mohler et al., 2012; Monazzam et al., 2014; Huss et al., 2015; Eyvazlou et al., 2016; Tettamanti et al., 2020), cognitive deficiency (Hutter et al., 2006), psychiatric symptoms (Silva et al., 2015) and microwave cataracts (Zaret, 1973). Thus they did not report a detailed description of the complete symptomatic picture of people associated with EMF exposure.

Surprisingly, only few studies have focused specifically on the description of the health symptoms in EHS self-reported patients. Most of these studies were also based on mail or web-based questionnaire and not on face-to-face questioning and examining patients. Such observational investigations concluded that symptoms are subjective, non-specific and not causally related to ELF, RF or MW EMF exposure (Levallois, 2002; Rösli, 2008; Rösli et al., 2010b; Baliatsas et al., 2014). However more recently studies allowing a more precise description of symptoms in such patients were conducted in Finland (Hagström et al., 2013) and in the Netherlands (van Dongen et al., 2014). In both studies, the percentage of women was higher in the EHS group than in the general population, suggesting some genetic susceptibility of these categories of patients, as reported in other studies including our own (Belpomme et al., 2015). In the Dutch study the number of symptoms was higher among people recruited by non-governmental organizations than in the general population (van Dongen et al., 2014), while in the Finnish study it was shown that the number of symptoms during the acute phase of EHS is higher than before its onset (Hagström et al., 2013). Table 2 summarizes all known major original published studies including our own reporting the symptomatic picture in EHS patients.

In fact, as emphasized by several scientists (Carpenter, 2015), the strongest evidence that EHS is a real syndrome similar to the microwave syndrome comes from the initial cases reported from 1980 to 2000 of acute high intensity exposure to MW EMF of healthy people, resulting in “prolonged illness” (Williams and Webb, 1980; Forman et al., 1982; Schilling, 1997, 2000; Reeves, 2000). Moreover, since it was shown that MCS is associated with EHS in near 25% of the EHS cases (Belpomme et al., 2015) and that both disorders are associated with inflammation, OS, possible BBB opening and brain neurotransmitter changes (Belpomme et al., 2015; Irigaray et al., 2018a; Belpomme and Irigaray,

**Table 2**

Major original published studies describing the symptomatic picture of EHS self-reported patients.

| Author                                    | Study types                                                   | Source/exposure                                                                                               | Total/evaluable cases                                                                                                                                                                        |
|-------------------------------------------|---------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Dodge, 1969 (USA)                         | Observation study                                             | MW                                                                                                            | 391 cases vs 100 controls                                                                                                                                                                    |
| Rea et al., 1991 (USA)                    | Provocation test                                              | 0.1 Hz–5 MHz EMF exposure                                                                                     | 25 patients vs sham and vs 25 healthy controls                                                                                                                                               |
| Bergqvist and Vogel 1997 (International)  | Nationwide questionnaire-based survey                         | General EMF exposure                                                                                          | 72 EHS patients                                                                                                                                                                              |
| Hillert et al., 2002 (Sweden)             | Population-based questionnaires                               | EMF, all types                                                                                                | 15.000 participants (general population), including 1.5% EHS patients                                                                                                                        |
| Navarro et al., 2003 (Spain)              | Questionnaire-based survey and EMF power density measurements | WC EMFs                                                                                                       | 101 persons close to MT base station                                                                                                                                                         |
| Oberfeld et al., 2004 (Spain)             | Questionnaire-based survey and EMF measurement                | WC EMFs                                                                                                       | 201 persons close to two GSM 900–1800 cellular phone base stations                                                                                                                           |
| Schreier et al., 2006 (Switzerland)       | telephone interviews cross-sectional study                    | 50/60 Hz EMF residential/personal exposure                                                                    | 2048 participants, including 5% (107) EHS patients                                                                                                                                           |
| Schüz et al., 2006 (Germany)              | Questionnaire-based survey via internet                       | EMF, all types including mobile phone use and MT base stations.                                               | 192 persons with health complaints, including 107 EHS patients                                                                                                                               |
| Röösli et al., 2010a, b (Switzerland)     | Population-based questionnaire and weekly measurements        | EMF, all types including MT base station proximity, mobile phone and cordless phone use and, W-LAN/WiFi.      | 1375 participants (general population), including 8% (130) EHS patients                                                                                                                      |
| Johansson et al., 2010 (Sweden)           | Questionnaire-based survey                                    | EMF, all types including domestic appliance and computer and mobile phone use                                 | 45 cases with mobile phone use and 71 EHS patients compared with a 106 population-based sample and 43 controls                                                                               |
| Kato and Johansson, 2012 (Japan)          | Questionnaire-base survey                                     | EMF, all types including medical device use, mobile phone and cordless use and proximity to MT base stations. | 75 EHS patients                                                                                                                                                                              |
| Hagström et al., 2013 (Finland)           | Questionnaire-based survey via internet                       | EMF, all types (selection of 50 electrical devices).                                                          | 194 EHS patients                                                                                                                                                                             |
| van Dongen et al., 2014 (The Netherlands) | Questionnaire-based survey via internet                       | EMF, all types                                                                                                | 188 people sensitive to EMF versus 937 people non-sensitive to EMF                                                                                                                           |
| Nordin et al., 2014 (Sweden)              | Questionnaire-based survey                                    | EMF, all types                                                                                                | 113 EHS patients versus 48 controls                                                                                                                                                          |
| Baliatsas et al., 2014 (The Netherlands)  | Questionnaire-based survey and electronic medical records     | EMF, all types including proximity to MT base stations, mobile phone use, domestic appliance and W-LAN/WiFi   | 5789 respondents including 514 (8.8%) cases with general environmental sensitivity and 202 cases (3.5%) with IEI-EMF (EHS) while the rest of respondents (5073 cases) were used as controls. |
| Belpomme and Irigaray, 2020 (France)      | Face-to-face physical examination                             | EMF, all types                                                                                                | 50 EHS, 50 EHS/MCS and 50 MCS people versus 50 apparently healthy people                                                                                                                     |

2020); it is believed that both EHS and MCS are objective somatic disorders, which cannot be claimed to originate from non-EMF-related psychologic or psychiatric cause, and neither result from a simple undefined and unproven functional impairment (Belpomme and Irigaray 2020, 2021; Belpomme et al., 2021)—although it cannot be excluded these disorders may occur in patients with some particular psychologic traits (Frick et al., 2002).

The purpose of provocation studies is to prove that EHS patients display acute symptoms at the time they are exposed (or after they are exposed) to man-made electric, magnetic and electromagnetic sources; whatever they are, i.e. ELF, RF or MW EMFs. As indicated above, the EHS-associated neurological symptoms are identical to those described in the MW syndrome which was considered at that time as evidently caused by MW EMF in exposed workers. A major difficulty here is that EHS patients are not only associated with hypersensitivity to low intensity anthropogenic EMFs, but due to their possible association with MCS, may also be sensitive to low concentration of multiple chemicals; so both environmental stressors could trigger clinical symptoms and pathological changes in these patients at weak or even very weak environmental EMF intensity or chemical concentration. Furthermore, against all standard medical practice, the clinical symptoms reported by the EHS patients have not been considered as medically assessed and recognized, but simply considered as “self-reported symptoms”, meaning they are not “functional symptoms”, as it is commonly used in medicine since Hippocrates. Hence they are not accepted as a valuable clinical descriptive tool to identify and diagnose EHS, due to their reported subjectivity and reported non-specificity. Moreover, it was claimed by WHO that EHS-associated symptoms differ from one patient to the other, a claim which is not confirmed by objective clinical observation analysis. In fact, as can be soundly deduced from any face-to-face questioning and physical examination of EHS patients, there is *a priori* no medical reason to dismiss the patients’ words, or to believe

they make up or mistake each time they attribute their symptoms to EMF exposure (Belpomme and Irigaray, 2020).

Many of the provocation studies performed in EHS patients were of insufficient methodological quality (Rubin et al. 2010, 2011). A major criticism as emphasized in the 2021 consensus report (Belpomme et al., 2021) is that these provocation tests were done before EHS had been objectively diagnosed using biomarkers and imaging techniques. This observation in addition to the flawed method used have resulted in negative findings. We thus consider *a priori* as scientifically unjustified to speculate that the electromagnetic claims of all the patients are unfounded and that their subjective symptomatic feeling could relate to some non-EMF psychosomatic or nocebo health effects (Belpomme et al., 2021; Belpomme and Irigaray, 2021). In Table 3 are depicted some of the unsuitable methodological issues of provocation tests having provided negative results.

An additional important reason for negative results in provocation studies is the fact that in cases of chronic suffering, the patients’ response to EMF exposure may be confused without clearly discriminating on/off or off/on field transition, especially when changes occur in a high rate with short-term field durations. In such cases a correct response to short time stimuli should not indeed be reasonably expected.

In fact not all provocation studies have provided negative results. Therefore, the apparently negative results could not preclude an absence of EMF trigger effects. Indeed, in well-designed provocation studies, ELF and/or WC pulsed RF or WC MW EMFs have been shown to trigger clinical and biological health effects in EHS patients. As indicated in Table 4, in such single- or double-blind provocation studies, various clinical and pathophysiological changes have been evidenced in these patients. Clinical effects include heart rate variability (HRV) and/or blood pressure variability (Havas et al., 2010; Havas, 2013; Koppel et al., 2018), altered pupillary light reflex (Rea et al., 1991), reduced visual perception (Trimmel and Schweiger, 1998), and abnormal

**Table 3**

Some unsuitable methodological issues in provocation tests of previously published studies having provided negative results (Belpomme et al., 2021).

|   |                                                                                                                 |                                                                                                               |
|---|-----------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------|
| 1 | Lack of precise inclusion criteria. No objective criteria based on molecular biomarkers and imaging techniques. | Röösli, 2008; Röösli et al., 2010b; Baliatsas et al., 2012; Schmiedchen et al., 2019                          |
| 2 | No clear consideration on medical anamnesis and degree of EHS severity.                                         | Baliatsas et al., 2012; Schmiedchen et al., 2019                                                              |
| 3 | No consideration of an association with MCS.                                                                    | Belpomme et al. 2015                                                                                          |
| 4 | No consideration that EHS patients are intolerant to specific man-made EMF frequencies.                         | Röösli, 2008; Röösli et al., 2010b; Baliatsas et al., 2012; Schmiedchen et al., 2019                          |
| 5 | Too short exposure duration.                                                                                    | Baliatsas et al., 2012; Eltiti et al., 2015                                                                   |
| 6 | Symptom recording made too early.                                                                               | Baliatsas et al., 2012; Schmiedchen et al., 2019                                                              |
| 7 | Endpoint criteria depending on subjective statements.                                                           | Röösli, 2008; Rubin et al., 2010, 2011; Baliatsas et al., 2012; Eltiti et al., 2015; Schmiedchen et al., 2019 |
| 8 | Possible EHS-associated psychological conditioning due to past suffering.                                       | Dieudonné, 2016                                                                                               |
| 9 | Possible significant EMF levels during sham exposure.                                                           | Alasdair, 2002                                                                                                |

movement during sleep (Mueller and Schierz, 2004), which all have been established by objective clinical evaluation. In addition, pathophysiological effects include altered electroencephalogram (EEG) during sleep (Arnetz et al., 2007; Lustenberger et al., 2013), altered electromyogram (EMG) after wireless local area network (WLAN) exposure (Tuengler and von Klitzing, 2013; von Klitzing, 2021), altered skin capillary blood flow (SCBF) (Tuengler and von Klitzing, 2013; Loos et al., 2013), and electric skin potential (ESP) and conductance changes (Tuengler and von Klitzing, 2013) – these also all allow objective measurements.

Moreover, in a single EHS case double-blind experiment, EMF-related symptomatic intolerance in comparison with sham-exposure has also been reported to be induced by off/on or on/off field transition, rather than by EMF uninterrupted exposure. As the authors state,

**Table 4**

Provocation tests performed in EHS patients using EMF exposure versus sham-exposure and/or comparison with healthy controls resulting in a positive causal link between EMF exposure and symptoms occurrence and/or pathophysiological changes.

| Study                                        | Endpoints                                      | Source            | Type of study                                            | EHS patients<br>Evaluable cases             | Results (effect of EMF exposure)                                                                                                                                                                                                                                                                 |
|----------------------------------------------|------------------------------------------------|-------------------|----------------------------------------------------------|---------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Rea et al., (1991)<br>(USA)                  | Pupillary light reflex                         | ELF (1–10 kHz)    | Double blind EMF v. sham provocation study               | 25 EHS patients' versus 25 healthy controls | 16/25 EHS patients consistently report symptoms in active, but not inactive conditions, compared with 0/25 healthy controls                                                                                                                                                                      |
| Trimmel and Schweiger (1998)<br>(Austria)    | Attention, perception and memory tests         | ELF (50 Hz)       | Double blind provocation study                           | 36 EHS versus 30 healthy controls           | Reduced performance of visual attention and perception by combining a 50 Hz magnetic field with acoustic noise exposure, compared to the effects of noise only.                                                                                                                                  |
| Mueller and Schierz (2004)<br>(Switzerland)  | Sleep disturbance                              | ELF (50 Hz)       | Double-blind cross-over provocation study                | 54 EHS cases                                | Cases moved away from area with maximum 50 Hz field intensity                                                                                                                                                                                                                                    |
| Arnetz et al., (2007)<br>(USA)               | Sleep EEG                                      | RF (884 MHz)      | Double blind case-control study compared to sham.        | 38 IEI-EMF and 31 healthy controls          | Exposure caused longer latency to deep sleep from sleep onset and reduced amount of cerebral slow wave                                                                                                                                                                                           |
| Mc Carty et al. 2011<br>(USA)                | symptomatic responses and EMF field perception | ELF (60 Hz)       | Single Blind provocation study, EMF versus sham exposure | A single female EHS case                    | In the first experiment, the EHS person reported somatic reactions with a significant difference with sham. In the second, she reported significantly more intense symptoms during exposure to a pulsed EMF in comparison with sham. In the third, she was not able to perceive EMF consciously. |
| Havas et al., 2010<br>(Canada)               | HRV, RBC clumping                              | RF (2.4 GHz)      | Single Blind provocation study EMF versus sham exposure  | 25 EHS self-reported patient                | 40% of EHS patients experienced some changes in their HRV during pulsed microwave exposure                                                                                                                                                                                                       |
| Tuengler and von Klitzing, 2013<br>(Germany) | HRV, capillary blood flow and SEP              | RF (Mobile phone) | Single Blind provocation study                           | Several types of EHS patients               | Modifications of biological parameters caused by EMF exposure                                                                                                                                                                                                                                    |
| Koppel et al., 2018<br>(Estonia)             | HRV                                            | ELF (50 Hz)       | Single Blind provocation study                           | 108 EHS patients                            | HRV significantly lower during EMF exposure than non-exposure.                                                                                                                                                                                                                                   |
| Von Klitzing, 2021<br>(Germany)              | ECG and EMG.                                   | RF (WiFi)         | Single Blind provocation study                           | 5 EHS patients                              | Modification of EMG caused by WLAN- exposure.                                                                                                                                                                                                                                                    |

this means that “the statistically reliable somatic reactions to subliminal EMF exposure were obtained under conditions that reasonably excluded the causative effect of any psychological process” (McCarty et al., 2011).

Such positive effects recorded by provocation tests have also been independently shown in two different earlier EHS case reports (Hocking and Westerman, 2002, 2003) and more recently in two studies showing the objective WC EMF effect on HRV in EHS patients in a double-blind provocation study (Havas et al., 2010) and more generally the effects of RF/MW EMF on the blood, the heart and the autonomic nervous system (Havas, 2013). Provocation studies using similar objective endpoints were also independently provided by the two German biophysicists Andreas Tuengler and Lebrecht von Klitzing, who considered that HRV, SCBF, ESP, and EMG recordings are suitable non-invasive methods to measure EHS in EHS patients (Tuengler and von Klitzing, 2013; von Klitzing, 2021). The same authors propose to combine the continuous measurements of HRV, SCBP and ESP overtime via electrocardiogram (ECG), Doppler meter and electrode matrix recordings respectively; before, during and after EMF versus sham-exposure. This method possibly allow the distinction of EHS patients from individuals suffering from other pathological conditions (Tuengler and von Klitzing, 2013).

As summarized in Table 4, objective abnormalities include the EHS-associated acute and reversible sympathetic and parasympathetic symptoms such as HRV and pupillary light reflex, and other acute neurological symptoms such as attention/memory loss and sleep disturbance, and above all objective biophysical cerebral and transient skin parameter changes, but not all symptoms are acute and reversible. In case of no treatment and no protective measures, chronic symptoms (such as loss of immediate and retrospective memory, mental confusion, insomnia, chronic fatigue, depressive tendency with possible suicidal ideation) may persist for a long time and even become irreversible, leading in some cases, to cerebral atrophy. Such evolution may occur in the case of chronic brain vascular insufficiency caused by persisting high resistance of the brain blood flow and low pulsatility in the cerebral middle arteries (Belpomme and Irigaray, 2020).

In fact, in EHS patients there seems to be a continuum from acute to

chronic symptoms, and from biological to health effects/disease; in case of no treatment and/or no efficient protection. We postulate two stages of EMF-related disease progression: first, where EMF-related biological effects may occur with a minimum of clinical symptoms; second, where pathophysiological changes and health symptoms predominate and lead to chronic disease. While the first step may be reversible, the second may be characterized by presumed pathological neurological lesions which may persist and be irreversible (see further).

Taking into account for all available scientific data we believe that present scientific knowledge strongly suggests that man-made EMF exposure can be causally involved in triggering harmful adverse clinical symptoms and noxious pathophysiological changes in EHS patients; and consequently that today's evidence of EMF-related multi-organic somatic effects dismisses the hypothesis of a causal psycho-pathological mechanism to account for the EHS-associated symptom occurrence.

#### 4. Search for electrohypersensitivity characterization

There remains persisting confusion between EHS, which was acknowledged by WHO (WHO, 2005; and IEI-EMF, which was proposed one year before, during the 2004 WHO-sponsored Prague meeting (Mild et al., 2006). EHS as indicated above is presently considered by WHO as a disability condition not proven to be causally related to EMF, and so not specifically subject to medical diagnosis, treatment and prevention; while IEI-EMF is defined as an idiopathic environmental intolerance condition possibly attributed to EMF.

We have proposed to define EHS as the intra-corporal acquisition of a pathological state of hypersensitivity to man-made EMFs in genetically or epigenetically predisposed EHS persons, as is the case for man-made chemicals in MCS patients (Belpomme et al., 2021). By contrast, IEI could be defined as the environmental intolerance to man-made EMFs, chemicals or other stressors, without the necessary acquisition of a state of hypersensitivity. More precisely, we proposed to define EHS clinically and biologically as a decrease in the physiological central nervous system (CNS)-associated EMF tolerance threshold, meaning that intolerance to EMF in EHS patients could occur for weak or even very weak EMF intensities, while intolerance to EMF in non-EHS people could occur for higher EMF intensities (Belpomme and Irigaray, 2021). We thus propose that designation of EHS be restricted to the presumable pathological intra-corporal acquisition of hypersensitivity to EMF, while IEI-EMF will be *stricto sensu* defined as presumable EMF-related environmental intolerance. A similar pathophysiological process involving a decrease in the CNS-associated chemical tolerance threshold could apply to MCS, a consideration that could result similarly in chemical intolerance for weak or even very weak concentrations of multiple environmental chemicals. Note that such a proposed pathophysiological definition, based on a decrease in the environmental tolerance threshold to better define EHS and MCS, is similar to that of toxicant-induced loss of tolerance proposed by Claudia S Miller (1999) who introduced this new concept of environmental sensitivity-related diseases.

While the present medical state-of-the-art must avoid any psychological causal interpretation for EHS occurrence and symptomatic development, there remains a first-order pending question: could the provocation tests prove hypersensitivity to man-made EMFs, i.e. that EHS patients are more sensitive to man-made EMFs than non-EHS healthy subjects; and could these patients detect the presence of ELF or WC RF/MW EMFs better than other persons? Relative to these two important question it was initially believed that using provocation tests in healthy people would show less or no responses under exposure to EMF in comparison with EHS patients (Wagner et al., 2000; Kleinogel et al., 2008; Valentini et al., 2010; Baliatsas et al., 2015). Similar results would be also expected in case-control studies (Landgrebe et al., 2008) or double blind provocation studies (Lowden et al., 2011); whereas EHS patients depending on the endpoint considered would exhibit typical responses during and/or after EMF-exposure. This is not the case. Contrary to previous supposition of none or fewer effects of man-made EMF

exposure in normal healthy individuals; many provocation studies, mostly using ELF and RF non-thermal man-made EMFs in healthy volunteers, have evidenced biological effects; while most studies in EHS patients were negative for the afore-mentioned reasons. The type of EMF/EMR used in provocation studies in healthy people is indicated in Table 5. These effects consist of decreased  $\beta$ -trace protein (prostaglandin D synthase) peripheral blood concentration (this molecule is an endogenous sleep promoting neurohormone) (Hardell et al., 2010), alterations of sleep EEG (Mann and Rösche, 1996; Schmid et al., 2012) and resting EEG (von Klitzing, 1995; Huber et al., 2002; Ghosn et al., 2015; Loughran et al., 2019), alteration of evoked electric potentials (Carrubba and Marino, 2008) and changes of the EEG alpha rhythm (Croft et al., 2008; Vecchio et al., 2012) and of the EEG slow beta, fast beta and gamma bands (Roggeveen et al., 2015). Such exposure to ELF or mostly to RF EMF (see Table 5) have also been shown in healthy subjects to alter the brain response during a memory task (Krause et al., 2000), to affect sleep dependent performance improvement in normal subjects (Lustenberger et al., 2013), to modify the 50 Hz exposure-induced human performance and psychophysiological parameters (Crasson et al., 1999), to induce annoyance and alter well-being (Zheng et al., 2015; Miller et al., 2019), to modify smells (Carlsson et al., 2005), and to influence cognitive performance (Verrender et al., 2016). In addition it has been reported that cell phone-associated WC EMF exposure decreases slow brain potentials at the central and temporo-parieto-occipital brain region (Freude et al., 1998), increases brain glucose metabolism activity (Volkow et al., 2011) and oxygen consumption at the frontal cortex (Curcio et al., 2009), alters non-thermal RFR-induced hemoglobin deoxygenation in cell-free preparations (Mousavy et al., 2009; Muehsam et al., 2013), influences electric properties of human blood measured by impedance spectroscopy (Sosa et al., 2005), increases blood viscosity (Tao and Huang, 2011), modifies brain vascularization (Huber et al., 2002; Aalto et al., 2006), alters blood pressure-associated baro-reflex activity (Braune et al., 1998), and induces vagal nerve stimulation at ECG and EEG (Burgess et al., 2016). In addition it has been shown that cell phone-induced HRV is dependent on breath, i.e. on the inspiration/expiration ratio (Béres et al., 2018). Most of these experimental studies in healthy people are summarized in Table 5, specifying the type of EMF/EMR exposure involved.

The hypothesis that EHS patients are really more sensitive to man-made EMF than healthy people, and that they could detect the presence of EMFs better than healthy people, is challenged by biological studies (Marková et al., 2005) as well as by epidemiological studies (Röösli, 2008) and provocation studies (Rubin et al., 2011); showing no evidence that short-term exposure to WC EMFs in EHS patients can cause self-reported symptoms, and that these patients could be able to detect ELF, RF or MW EMF better than healthy subjects.

Considering the above reported EMF-induced positive effects in healthy people, it will be extremely difficult to scientifically demonstrate the specific EMF-related hypersensitivity state in EHS patients, i.e. their sensitivity to lower intensity EMFs, using comparative methods. Therefore, research on hypersensitivity to EMFs using such clinical approach in EHS patients may remain an open question for a long time. Although the toxic pathophysiological role of EMF has been ascribed in different animal and human studies, this role has still not been studied specifically for EHS.

#### 5. Search for etiology

The uncertain results of many provocation tests performed in EHS self-reported patients and their misinterpretation have resulted in postulating some placebo effects; accounting for the great confusion existing presently between researchers within the scientific and medical community and consequently within the international and national medical, sanitary and societal institutions. A big mistake is that the negative results provided by these provocation studies have been

**Table 5**

Double or single blind provocation studies or observational studies resulting in positive EMF-associated causal link in healthy volunteers.

| Authors, Year, Country            | Endpoints                                                                                          | Type of study                                                                                                               | Evaluable cases                                                     | Results (effect of EMF exposure)                                                                                                                                                                                    |
|-----------------------------------|----------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| von Klitzing L. 1995 (Germany)    | Changes in resting EEG                                                                             | Observational study involving low frequency (217 Hz) exposure                                                               | 17 healthy students                                                 | Alteration in the range of alpha-activity during and after exposure for some hours                                                                                                                                  |
| Mann and Roschke, 1996 (Germany)  | Changes in sleep EEG                                                                               | Single blind study involving RFR (900 MHz) exposure                                                                         | 24 healthy male volunteers                                          | Temporal pattern of cortisol secretion differs between placebo and night exposure                                                                                                                                   |
| Braune et al., 1998 (Germany)     | Blood pressure (BP), heart rate, capillary perfusion, and subjective well-being                    | Single-blind placebo-controlled study involving RFR (900 MHz) exposure                                                      | 7 healthy volunteers                                                | BP associated baro-reflex with activity alteration                                                                                                                                                                  |
| Freude et al., 1998 (Germany)     | Slow brain potentials (SBP)                                                                        | Single blind study involving RFR (916.2 MHz) exposure                                                                       | 16 healthy young people                                             | significant decrease of SBP in central and temporo-parieto-occipital brain regions                                                                                                                                  |
| Crasson et al., 1999 (Belgium)    | Changes in event-related potentials (ERP) and EEG/psychophysiological and psychological behavior   | Two double blind experimental studies involving 50 Hz exposure and sham                                                     | 21 healthy male volunteers                                          | Low level 50 Hz MF may have a slight influence on ERP and reaction time under circumstances of sustained attention.                                                                                                 |
| Krause C.M. 2000 (Finland)        | Changes in EEG (during a memory task)                                                              | Single blind study involving RFR (902 MHz) exposure                                                                         | 16 healthy volunteers                                               | RFR modifies the brain responses                                                                                                                                                                                    |
| Croft et al., 2002 (Australia)    | effects of active mobile phone (MP) on the neurological system                                     | Single blind cross-over study involving RFR (900 MHz) exposure                                                              | 24 healthy volunteers                                               | MP exposure affects brain functioning                                                                                                                                                                               |
| Huber et al., 2002 (Switzerland)  | Effect of EMF on waking regional cerebral blood flow (rCBF) and on waking and sleep EEG in humans. | Double blind study involving two types of RFR (a 'base-station-like' and a 'handset-like' signal) vs. sham control exposure | 16 healthy young male right-handed subject                          | Pulse EMF increases waking rCBF and pulse modulation of EMF is necessary to induce waking and sleep EEG changes                                                                                                     |
| Curcio et al., 2005 (Italia)      | Effects of GSM on the neurological system:                                                         | RFR (902.4 MHz) exposure                                                                                                    | 20 healthy volunteers                                               | EMF affects normal brain functioning                                                                                                                                                                                |
| Carlsson et al., (2005) (Sweden)  | Annoyance related to electrical and chemical factors in a Swedish general population               | Cross-sectional study involving different electrical equipment.                                                             | 13,604 subjects, representative of the population of Scania, Sweden | Connection between environmental annoyance, well-being and functional capacity                                                                                                                                      |
| Huber al., 2005 (Switzerland)     | Effect of EMF on waking regional cerebral blood flow (rCBF)                                        | Double blind study involving two types of RFR (a 'base-station-like' and a 'handset-like' signal) vs. sham control exposure | 12 healthy young male subjects                                      | Only 'handset-like' RFR exposure affected rCBF                                                                                                                                                                      |
| Aalto et al., 2006 (Finland)      | Effects of an active mobile phone on rCBF                                                          | Double-blind, counterbalanced study design with subjects performing a computer-controlled verbal working memory task        | 12 healthy volunteers                                               | EMF emitted by a commercial mobile phone affects rCBF in humans                                                                                                                                                     |
| Croft et al., 2008 (Australia)    | Effects of MP on the neurological resting system                                                   | Double blind cross-over study. RFR (895 MHz) exposure versus sham.                                                          | 120 healthy volunteers                                              | Alpha power enhancement during MP exposure                                                                                                                                                                          |
| Carrubba and Marino, 2008 (USA)   | Evoked brain electrical potentials, EEG normal humans, and patients with epilepsy                  | Review on different normal human studies                                                                                    | Different normal human studies                                      | Changes in brain activity                                                                                                                                                                                           |
| Curcio et al., 2009 (Italy)       | Oxygenation of the frontal cortex by functional near-IR spectroscopy (fNIRS)                       | Double blind Case-control study of GSM signal (902.4 MHz) compared to sham.                                                 | 31 healthy students                                                 | Slight influence in frontal cortex                                                                                                                                                                                  |
| Moussavy et al., 2009 (Iran)      | Structure and function of hemoglobin                                                               | Experimental study involving RFR (910 MHz and 940 MHz) exposure                                                             | Human adult hemoglobin prepared from human RBC of healthy donors.   | MP electromagnetic fields decreases oxygen affinity and modifies tertiary structure of hemoglobin depending on field intensity and time of exposure.                                                                |
| Hardell et al., 2010 (Sweden)     | Effect of MP and/or cordless phone on $\beta$ -trace protein blood concentration                   | Observational study involving RFR (MP and cordless phone)                                                                   | 62 health volunteers                                                | Long term wireless phone use decreases $\beta$ -trace protein                                                                                                                                                       |
| Carrubba et al., 2010 (USA)       | Effects of MP (217 Hz) on the neurological system:                                                 | Double blind study                                                                                                          | 20 healthy volunteers                                               | MP trigger evoked potentials at the frequency of 217 Hz during ordinary MP use.                                                                                                                                     |
| Lowden et al., 2011 (Sweden)      | Sleep EEG                                                                                          | RFR (884 MHz) exposure versus sham double blind study                                                                       | 48 healthy volunteers                                               | RFR exposure increases alpha range in sleep EEG                                                                                                                                                                     |
| Volkow et al., 2011 (USA)         | Brain glucose metabolism (PET-scan)                                                                | Single blind study involving 50 min cell phone (837.8 MHz) exposure                                                         | 47 healthy participants                                             | Increased brain glucose metabolism in the region closest to the antenna                                                                                                                                             |
| Tao and Huang 2011 (USA)          | Blood viscosity                                                                                    | Experimental study involving 1.3 T magnetic pulse to a small sample of blood                                                | Human blood from healthy donors                                     | After 1 min of exposure blood viscosity is reduced by 33%                                                                                                                                                           |
| Vecchio et al., 2012 (Italy)      | Changes in GSM event-related desynchronisation (ERD) at resting EEG                                | Placebo controlled double blind study involving RFR (902.4 MHz) exposure                                                    | 11 healthy volunteers                                               | The peak amplitude of $\alpha$ ERD and the reaction time to go stimuli are modulated by the effect on the cortical activity                                                                                         |
| Schmid et al., 2012 (Switzerland) | Resting EEG and polysomnography cognitive/behavioral endpoints                                     | Double blind cross-over study involving RFR (900 MHz) exposure                                                              | 30 young healthy men                                                | pulse-modulated RFR alter brain functioning                                                                                                                                                                         |
| Muehsam et al., 2013 (USA)        | Structure and function of hemoglobin                                                               | Experimental study involving a pulse-modulated RFR (27.12 MHz) or a static magnetic field exposure                          | Human adult hemoglobin prepared from human RBC of healthy donors.   | Exposure for 10–30 min to either pulse-modulated radiofrequency or static magnetic field increased the rate of deoxygenation of hemoglobin occurring several minutes to several hours after the end of EMF exposure |

(continued on next page)

Table 5 (continued)

| Authors, Year, Country                  | Endpoints                                                              | Type of study                                                                         | Evaluable cases                       | Results (effect of EMF exposure)                                                                                                                      |
|-----------------------------------------|------------------------------------------------------------------------|---------------------------------------------------------------------------------------|---------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------|
| Lustenberger et al., 2013 (Switzerland) | Brain activity during sleep EEG                                        | Double blind cross-over study involving RFR (900 MHz) exposure                        | 16 healthy male people                | RFR affect ongoing brain activity during sleep                                                                                                        |
| Ghosh et al., 2015 (France)             | Changes in resting EEG effects of GSM on the neurological system       | Double blind Case-control study compared with sham involving RFR (900 MHz) exposure.  | 26 healthy volunteers                 | During exposure and post-exposure, the alpha band power is significantly decreased with closed eyes compared to sham.                                 |
| Roggeveen et al., 2015 (UK)             | Changes in resting EEG                                                 | Single blind, cross-over study involving RFR (1.9291–1.9397 GHz) exposure             | 31 young female                       | All brain waves except delta change significantly due to exposure of the ear, in comparison to sham, with stronger effects with ipsilateral exposure. |
| Burgess A.P. et al., 2016 (UK)          | Resting EEG and ECG (HRV)                                              | Blinded randomized provocation study with a standardized TETRA signal versus sham     | 164 police officers and 60 volunteers | vagal nerve stimulation at ECG and EEG                                                                                                                |
| Verrender et al., 2016 (Australia)      | Visual discrimination task and modified Sternberg working memory task, | Double blind cross-over study involving pulse modulated RFR (PMRF) (920 MHz) exposure | 36 healthy volunteers                 | Cognitive performance is faster relative to sham in a working memory task during PMRF exposure.                                                       |
| Béres et al. 2018 (Hungary)             | Heart rate asymmetry (HRA) and HRV parameters using repeated-measures  | Double-blind crossover study involving RFR (1800 MHz) exposure                        | 20 healthy volunteers                 | Increased HRV under 1:1 breathing and RFR exposure                                                                                                    |
| Loughran et al., 2019 (Australia)       | Changes in resting EEG                                                 | Double blind cross-over study involving RFR (920 MHz) exposure versus sham            | 36 healthy volunteers                 | Alpha activity increases during high exposure condition compared to sham                                                                              |

interpreted not to arise from their incorrect methodological practice (Blackman, 2009; Schmiedchen et al., 2019; Belpomme et al., 2021) but rather from some nocebo effect, considering EHS as a psychological disease (Rubin et al. 2010, 2011). Indeed the so called nocebo effect is at best a hypothesis that needs to be confirmed by suitable experimental studies (Belpomme et al., 2021; Belpomme and Irigaray, 2021). This has not occurred. To the contrary, on the basis on a limited number of interviews of EHS patients, it has been suspected that the psycho-social behavior associated with EHS in these patients is secondary to disease occurrence and suffering, a consequence and not a cause of EHS (Dieudonné, 2016). Moreover, the molecular (Belpomme et al., 2015; Irigaray et al., 2018a; Belpomme and Irigaray, 2020) and radiological abnormalities (Heuser and Heuser, 2017; Irigaray et al., 2018b; Greco, 2020) that have been detected in EHS patients demonstrate that EHS is a neurological somatic disease not a psychological disease. Similarly, MCS has not only been shown to be associated with increased sensitivity to multiple chemicals, but also to be caused by some initial acute or sub-acute toxic episodes triggered by environmental chemicals – mostly synthetic – in genetically susceptible hosts (Bartha et al., 1999). Therefore neither MCS nor EHS can be considered to be of psychological origin. Also, EHS may be characterized not only as a specific state of intolerance to low intensity EMFs, but also as caused by previous excessive EMF exposure. This critical interpretation was initially provided by David Carpenter by analyzing the microwave syndrome (Carpenter, 2014, 2015). This concept was more recently developed in a review analyzing the EHS underlying mechanisms involving EMF exposure by Y. Stein and I.G. Udasin (2020).

In Table 6 the prevalence expressed in percentages of EHS people relative to the overall population is estimated to range from 0.7% to 13.3%, mainly affecting on average 3%–5% of the population in many different worldwide area or countries, meaning that millions of people may in fact be affected by man-made EMF intolerance, and often by EHS. Similar worldwide figures may account for MCS (Genuis, 2010).

From the analysis of our data and those of the scientific literature, we now consider several strong and convincing arguments that prove EHS is caused by non-thermal anthropogenic EMF exposure.

1. EHS cannot be considered to originate from a nocebo effect i.e. be a psychiatric disease; due to the findings showing its association with somatic abnormalities such as low grade inflammation, OS, and consequent disruption/opening BBB as well as in some cases with anti-myelin Po autoimmune response (Belpomme et al.,

2015; Belpomme and Irigaray, 2021). EHS should be therefore considered a somatic disease. In addition we have shown it is associated approximately in 25% of the cases with MCS which is already considered as a somatic disorder (Belpomme and Irigaray, 2021). Moreover EHS is an increasing worldwide plague, hence it is reasonably expected not to be a nocebo disease.

2. EHS occurrence has appeared subsequently to artificial electromagnetic environmental pollution with a seemingly progressive increasing prevalence since the use of WC technologies (Bandara and Carpenter, 2018).
3. As indicated in Table 6 intolerance to EMF exposure including EHS occurrence is not restricted to some regional areas or to countries, but is a worldwide plague with pandemic extension, as is the case for the worldwide expansion of the EMF emitting technologies (Hallberg and Oberfeld, 2006; Bandara and Carpenter, 2018).
4. There are many independent provocation studies proving that ELF/RF/MW EMF can biologically damage the organism and are noxious agents in healthy people (see Table 5); while due to the use of incorrect methodology (see Table 4) in EHS suffering patients, there is a limited number of studies showing pathophysiological changes and symptoms induction. Therefore negative provocation studies definitely cannot exclude a causal role of EMFs in EHS patients.
5. Several main EHS-associated symptoms such as sleep disturbance (Davis, 1997), depressive tendency (Poole et al., 1993; Verkasalo et al., 1997) and suicide risk (Perry et al., 1981; Johnston, 2008) have been shown in independent epidemiological studies to result from dose-dependent EMF exposure, implying that excessive EMF exposure is the cause of these characteristic EHS-associated symptoms (Perry et al., 1981; Poole et al., 1993; Davis, 1997; Verkasalo et al., 1997; Johnston, 2008).
6. As previously reported many EHS patients are characterized by possible low grade inflammation, nitroso-oxidative stress, BBB disruption/opening and brain neurotransmitter changes (Belpomme et al. 2015, 2018; Irigaray et al., 2018a; Belpomme and Irigaray, 2020); all which have been shown in laboratory animals by different independent studies to be caused by man-made EMF exposure (Salford et al. 1994, 2003; Cao et al., 2000; Eberhardt et al., 2008; Nittby et al., 2009; Yang et al., 2012; Aboul Ezz et al., 2013; Megha et al. 2015a, 2015b; Saili et al., 2015; Hu et al., 2021).

**Table 6**

Estimated prevalence of people with self-reported intolerance to EMF and/or EHS in different countries.

| Author, Year, Country                  | Year of results | Sample Size                 | People Contribution Rate (%)** | Estimated % of People with EHS |
|----------------------------------------|-----------------|-----------------------------|--------------------------------|--------------------------------|
| Hillert et al. (2002), Sweden          | 1997            | 15,000 (19–80) <sup>a</sup> | 73                             | 1.5                            |
| Palmquist et al. (2014), Sweden        | 2010            | 3406                        | 40                             | 2.7                            |
| Schreier et al. (2006), Switzerland    | 2004            | 2048 (>14) <sup>a</sup>     | 55.1                           | 5                              |
| Röösli et al., 2010a, Switzerland      | 2008            | 1122 (30–60) <sup>a</sup>   | 37                             | 8.6                            |
| Röösli et al., 2010b, Switzerland      | 2009            | 1122 (30–60) <sup>a</sup>   | 37                             | 7.7                            |
| Blettner et al. (2009), Germany        | 2004            | 30,047                      | 58.6                           | 10.3                           |
| Kowall et al. (2012), Germany          | 2004            | 30,047                      | 58.4                           | 8.7                            |
| Kowall et al. (2012), Germany          | 2006            | 30,047                      | 58.4                           | 7.2                            |
| Levallois et al. (2002), USA           | 1998            | 2072                        | 58.3                           | 3.2                            |
| Korpinen and Pääkkönen, 2009, Finland  | 2002            | 6121                        | 40.8                           | 0.7                            |
| Eltiti et al. (2007), UK               | 2005            | 3633                        | 18.2                           | 4                              |
| Meg Tseng et al. (2011), Taiwan        | 2007            | 1251                        | 11.5                           | 13.3                           |
| Schröttner and Leitgeb (2008), Austria | 2008            | 460                         | 88                             | 3.5                            |
| Furubayashi et al. (2009), Japan       | 2007            | 2472                        | 62.3                           | 1.2                            |
| Baliatsas et al. (2014), Netherlands   | 2011            | 5789                        | 39.6                           | 3.5                            |
| van Dongen et al., 2014, Netherlands   | Before 2013     | 1009                        | 60                             | 7                              |

<sup>a</sup> When provided age of included patients is indicated in brackets.

\*\*Contribution rate is the percentage of people having answered positively to the survey.

7. Most EHS patients present in their past medical history excessive exposure to WC RF/MW EMFs, and/or ELF EMFs, confirming that exposure to anthropogenic EMF may be a main plausible causal factor in inducing EHS (Belpomme and Irigaray, 2020).

8. Many independent *in vitro* and *in vivo* studies demonstrate that man-made EMFs can interact with endogenous physiological electric fields which control cellular biological functions in normal organism (Weisenseel, 1983; Nuccitelli, 1988, 2000; Borgens, 1988; Blanchard and Blackman, 1994; Shi and Borgens, 1995; McCaig and Zhao, 1997; McCaig et al., 2005; Yao et al., 2009; Del Giudice et al., 2011; Funk, 2015). When applied to the whole human organism, man-made EMFs distort the physiological endogenous EMFs. They also distort the corresponding cellular functions which results in adverse biological/health effects via EMF/tissue interaction at a molecular level (Blank, 2005; Vander Vorst et al., 2006). This is particularly the case for human brain, heart and muscles all being involved biologically and symptomatology in EHS, a finding confirming the multi-target causing role of man-made EMF-exposure (Frey, 1993; Vander Vorst et al., 2006).

9. It has been shown that man-made EMFs and their corresponding EMR are completely polarized and coherent, and thus differ physically from natural EMF/EMRs which are non-polarized. This key-difference may account for their harmful and toxic effects on biomolecules, cells and tissues, in contrast to natural EMFs, which are necessary for life (Panagopoulos et al., 2015a; Panagopoulos, 2017, 2019, 2021).

10. The pathophysiological mechanism by which polarized and coherent (man-made) EMFs may cause neurotoxic effects is now evidenced. Many *in vitro* and *in vivo* animal (Bas et al., 2009; Sonmez et al., 2010; Yang et al., 2012; Aldad et al., 2012; Deshmukh et al., 2013; Balassa et al., 2013; Furtado-Filho et al., 2015; Megha et al., 2015a; Zhang et al., 2015; Odaci et al., 2016; Sirav and Seyhan, 2016), and human studies (Gandhi et al., 1996; Cardis et al., 2008; Dasdag et al., 2012; Belpomme et al., 2018) evidence the neurological and mainly brain noxious effects of man-made non-thermal or micro-thermal EMFs.

11. At the molecular level it has been shown that non- or micro-thermal low-intensity/long duration EMF exposure act directly on DNA, not only by inducing DNA strand breaks or DNA fragmentation (Lai and Singh, 1995, 2004; Phillips et al., 2009; Panagopoulos, 2019; Lai, 2021), but also by inducing chromosome alteration (Sekeroglu et al., 2012, 2013) and chromatin modification (Belyaev and Kravchenko, 1994; Belyaev, 2005). In addition following genetic damage (Lai, 2021 appendix 1 and 2) and/or epigenetic changes (Blank and Goodman, 1999; Belyaev, 2005; Belyaev et al., 2006; Leone et al., 2014; Dasdag et al., 2015a; Dasdag et al., 2015b), EMF exposure could induce gene regulation changes (Lai, 2021 appendix 3) and protein misfolding (Millenbaugh et al., 2008). In fact, multiple cell targets following external application of EMF – mostly RFR and MW EMF – to the whole organism should be considered in different tissues including the brain. It is still unclear whether these different genetic and/or epigenetic mechanisms are involved in EHS genesis, but as shown in many studies, cell free radicals production following ELF or RF EMF exposure (Lai 2019) may take part in these alterations. We have shown that in 80% of the cases of EHS patients EHS is associated with the production of reactive oxygen species (ROS) and/or reactive nitrogen species (RNS) free radicals, suggesting that EMFs could be indirectly involved in EHS genesis (Irigaray et al., 2018a).

Furthermore, it has been shown that EMFs can interact directly with DNA in a specific magnetic field responsive domain in the HSP70 promoter to induce rapid synthesis of heat-shock proteins, a finding which can account for the anti-inflammatory response reported to occur in healthy people (Lin et al. 1999, 2001; Blank and Goodman, 1999, 2011; Blank, 2005); a result we have also shown to occur in EHS patients (Belpomme et al., 2015).

12. All these different findings clearly argue for a causal role of EMF in inducing EHS directly or indirectly via ROS and/or RNS. Although EMF exposure appears to be the main cause of EHS and can explain the pathophysiological change and the symptomatic occurrence, the specific mechanism of EHS genesis, i.e. the occurrence of a decrease in the EMF intolerance threshold is still hypothetical (see further). In addition, in some EHS cases MCS may precede the occurrence of EHS. Thus we have hypothesized that chemicals may also be implicated as causing agents in EHS genesis in a limited number of cases (11%) (Belpomme and Irigaray, 2020). Additionally in conjunction with the causal role of EMF and/or chemicals there may be some independent risk factors associated with EHS genesis, such as a preexisting depression, a psychiatric comorbidity (Meg Tseng et al., 2011), a previous brain trauma, a possible acquired immunosuppression-associated opportunistic infection, or a congenital malformation; which could further the EMF- and/or chemical-related EHS genesis in genetically and/or epigenetically predisposed individuals. Future research must focus on these different risk

factors with appropriate epidemiological studies and suitable bioclinical methods.

## 6. Hypothetical biophysical mechanisms specifically involved in EHS genesis

There are some further indications supporting the hypothesis of a particular biophysical mechanism, accounting specifically for a causal role of EMF in inducing hypersensitivity:

- (a) due to the presence of electromagnetic receptors, as in bacteria and many animals, humans are all sensitive to EMFs, but normally not hypersensitive. Such receptors have been identified as “cryptochroms” in animal retina (Gegear et al., 2010; Grehl et al., 2016) and as “magnetosomes” in the human brain (particularly in the hippocampus) and in the meninges (Kirschvink et al., 1992a; Dunn et al., 1995; Maher et al., 2016). Magnetosomes are located mainly in areas thought to correspond to the observed EHS-associated pathophysiological abnormalities and clinical symptoms (hippocampus and meninges) in EHS patients. These latter receptors have been shown to contain ferrous magnetite (grainite) and maghemite crystals (Kirschvink et al., 1992a) which have been thought to sense EMFs. Moreover, biogenic magnetite has been shown to be associated with ferromagnetic resonance and to absorb EMFs, hence it can be a mechanism capable of producing some biological response under the influence of EMF (Kirschvink et al., 1992b; Johnsen and Lohmann, 2005). Since these receptors are basically constituted of minerals they are thought to sense not only natural ELF, i.e. the Earth’s magnetic field but also man-made polarized static ELF EMF and man-made ELF-associated RFR. Humans may have indeed a geomagnetic sensory neurologic system as do many other animals. But most of them are not consciously aware of the Earth’s magnetic field that is encountered in everyday life (Wang et al., 2019). Possibly they have lost this shared magnetic sensory system due to the development of some hypothetical adaptive protection systems. The alteration (or destruction) of this putative anti-EMF adaptive neurologic system by excessive man-made EMF exposure (see further) may explain occurrence of hyper-sensitivity to EMF by restoring the remnant primordial sensing effect of magnetosomes. Restoration of other hypothetical EMF sensing receptors might be involved to account for the particular state of EHS.
- (b) At a molecular level it has been theorized that the voltage-gated ion channels (VGICs) in cell membranes could be a possible target for polarized and coherent (man-made) EMFs (Bawin and Adey, 1976; Liburdy, 1992; Waliczek, 1992; Balcavage et al., 1996; Panagopoulos et al., 2002, 2015b, 2021). It has been proposed that biogenic magnetite, under the influence of EMF can open such VGICs (Kirschvink et al., 1992b; Johnsen and Lohmann, 2005). But the VGICs physicochemical process which mainly involves calcium ions (Bawin and Adey 1976; Liburdy, 1992; Waliczek, 1992; Pall, 2013), has been thought to be applied to all cells in the organism. Therefore, it cannot explain the unique sensing mechanism/effect of EHS and the particular EHS-associated pathophysiological changes observed in the CNS; specifically in the hippocampus and the meninges. Other EMF-induced mechanisms/effects may be involved;
- (c) It has been shown in laboratory animals that EMFs and/or chemicals can particularly damage neurons (Frey, 1993; Redmayne and Johansson, 2014; Megha et al. 2015a, 2015b), and change the neurotransmitter and synapse-related protein concentrations particularly in the hippocampus (Bas et al., 2009; Leone et al., 2014; Teimori et al., 2016; Tan et al., 2019). Moreover, neurons are more vulnerable to EMF-induced apoptosis than other cells in the organism (Salford et al., 2003;

Joubert et al., 2008; Sonmez et al., 2010; Zuo et al., 2014; Odaci et al., 2016; Eghlidospour et al., 2017). Since as previously defined EHS appears clinically to be an acquired and persisting state, our hypothesis is that man-made EMFs and/or marketed chemicals in EHS patients may have permanently altered or destroyed neurons of the adaptive protective system, and their neuronal circuits in the brain, possibly in the hippocampus (Belpomme and Irigaray, 2020). This is a path for further biophysical and pathophysiological research efforts in order to better characterize (hyper)sensitivity of EHS and/or MCS, to eventually validate our proposed hypothesis via further specific CNS neurological investigations.

## 7. Discussion

By using several biomarkers in the peripheral blood and urine, and suitable cerebral imaging techniques (Irigaray et al., 2018b; Belpomme and Irigaray, 2021), we have previously evidenced that EHS is a brain pathological disorder which can be objectively diagnosed and treated. Moreover, it has been shown that, although they differ in their etiology and pathogenesis, both EHS and MCS share a similar clinical and biological signature, so they must be considered medically as parts of a particular unique environmental intolerance-related neurological syndrome (Belpomme et al., 2015). This is what many scientists recently agreed to in a scientific consensus report stipulating the critical role of biomarkers and imaging to study EHS (Belpomme et al., 2021). Our finding on EHS mainly based on the use of biomarkers and suitable imaging techniques must however be confirmed by other studies. But we show here that the present research progress results in the acknowledgment of EHS as a real pathological disorder caused by EMF exposure. Indeed further research efforts should be made to prove definitely the causal role of EMFs in triggering EHS-associated symptoms and EHS genesis itself. However, the different and independent data that we have provided fulfill the causation criteria proposed by WHO (WHO, 2006) because a) they include a dose-response effect of the main EHS-associated symptoms in epidemiological studies, b) they testify that the biological changes of *in vitro* and *in vivo* laboratory animals exposed to man-made EMFs are similar to what is observed in EHS patients, c) they also evidence an EHS-associated non-thermal or micro-thermal pathophysiological mechanism accounting for symptom occurrence, and d) above all they fully obey the general scientific principles used by different independent research teams. The data therefore supports the role of man-made EMFs as a causal agent of EHS. In addition, it is clearly demonstrated in different independent studies using provocation tests, that EMFs are noxious for healthy people. Consequently, there are sufficient established facts to strongly recommend protective measures against the present man-made electromagnetic pollution, using the precautionary principle to protect in particular pregnant women, infants, children, teenagers and young adults in all countries worldwide.

Given the seven billion people worldwide – most using cordless phones and/or mobile phones, Wi-Fi, and other wireless devices – and given the present and future development of 5G (Hardell and Nyberg, 2020; Hardell and Carlberg, 2020; Pall, 2021), it is expected that the prevalence of EMF intolerance and EHS will significantly increase worldwide in the next few years. However, because the figures indicated in Table 6 are estimations based on no objective criteria for identifying EHS (Hallberg and Oberfeld, 2006), we believe these data require confirmation by more objective evaluations. Although the reported EHS prevalence figures are only estimations, it is expected that EMF intolerance and EHS prevalence will continue to grow, in as much as the manufacturers of WC technologies and chemical industries will continue developing their products.

As reported in this overview, since the 2005 and 2014 WHO official publications, much progress has been made in the identification and understanding of EHS (and MCS) as pathological disorders and the bioclinical health effects of man-made EMFs and/or chemicals on the

organism. But EHS and MCS have still not been adequately acknowledged by WHO. The non-thermal or micro-thermal health effects of man-made EMF exposure evidenced in animals as well as in humans and their physico-chemical mechanisms of action (Pall, 2013; Yakymenko et al., 2016; Belpomme and Irigaray, 2020; Panagopoulos et al., 2021) should be considered by WHO. Contrary to the unrealistic claims by the International Commission on Non-Ionizing Radiation Protection (ICNIRP), who still denies the existence of non- or micro-thermal biological and toxic health effects of man-made EMFs (ICNIRP, 1998, 2010, 2020), we emphasize again that critical research progress has recently been made, making non- and micro-thermal EMF effects today a common acknowledgment among scientists and civil society people, as testified by many international scientific appeals calling for a reasonable limitation of electromagnetic pollution and the deletion or a *moratorium* of 5G development (Hardell and Nyberg, 2020; Hardell and Carlberg, 2020; Pall, 2021). Indeed, it appears that the non- or micro-thermal EMF-related toxic health effects (in addition to the MCS-related environmental effects) are the cause of EHS pathogenesis and etiology, as is also a possible cause of cancer (Hardell et al., 1995; IARC, 2002; Belpomme et al., 2007; IARC, 2013; Hardell et al., 2013).

Furthermore, the health care needs of people with environmental sensitivities such as EHS or MCS should be determined and developed in the present socioeconomic environment and medical challenge (Gibson et al., 2015).

Today's level of scientific knowledge engenders a great ethical responsibility of scientists and governments and of national and international health bodies to uncover the adverse health effects of the increasing man-made EMF exposure and warn on the emerging and growing worldwide EHS and MCS global plagues. This means that suitable public health measures must urgently be taken to recognize EHS and MCS as new pathologies and decrease EMF-exposure.

We therefore strongly ask WHO to add EHS and MCS in the future versions of the WHO International Classification of Diseases on the basis on their clinical and pathophysiological identification, just as has already been done for other recognized diseases.

## Funding

This research was funded by the European Cancer and Environment Research Institute (ECERI) grant number 0001092020.

## Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## References

- Aalto, S., Haarala, C., Brück, A., Sipilä, H., Hämäläinen, H., Rinne, J.O., 2006. Mobile phone affects cerebral blood flow in humans. *J. Cerebr. Blood Flow Metabol.* 26, 885–890.
- Aboul Ezz, H.S., Khadrawy, Y.A., Ahmed, N.A., Radwan, N.M., El Bakry, M.M., 2013. The effect of pulsed electromagnetic radiation from mobile phone on the levels of monoamine neurotransmitters in four different areas of rat brain. *Eur. Rev. Med. Pharmacol. Sci.* 17, 1782–1788.
- Alasdair, P., 2002. Peer Review and Quality of Science. Comments Posted at 07/08/2006 on Powerwatch Website. <https://www.powerwatch.org.uk/columns/aphilips/index.asp>.
- Aldad, T.S., Gan, G., Gao, X.B., Taylor, H.S., 2012. Fetal radiofrequency radiation exposure from 800–1900 mhz-rated cellular telephones affects neurodevelopment and behavior in mice. *Sci. Rep.* 2, 312.
- Arnetz, B.B., Akerstedt, T., Hillert, L., Lowden, A., Kuster, N., Wiholm, C., 2007. The effects of 884 MHz GSM wireless communication signals on self-reported symptom and sleep (EEG). An experimental provocation study. *PIERS Online* 3, 1148–1150.
- Auvinen, A., Feychting, M., Ahlbom, A., et al., 2019. Headache, tinnitus and hearing loss in the international cohort study of mobile phone use and health (COSMOS) in Sweden and Finland. *Int. J. Epidemiol.* 48, 1567–1579.
- Balassa, T., Varró, P., Elek, S., Drozdovszky, O., Szemerszky, R., Világi, I., Bárdos, G., 2013. Changes in synaptic efficacy in rat brain slices following extremely low-frequency magnetic field exposure at embryonic and early postnatal age. *Int. J. Dev. Neurosci.* 31, 724–730.
- Balcavage, W.X., Alvager, T., Swez, J., Goff, C.W., Fox, M.T., Abdullyava, S., King, M.W., 1996. A mechanism for action of extremely low frequency electromagnetic fields on biological systems. *Biochem. Biophys. Res. Commun.* 222, 374–378.
- Baliatsas, C., Bolte, J., Yzermans, J., Kelfkens, G., Hooiveld, M., Lebre, E., van Kamp, I., 2015. Actual and perceived exposure to electromagnetic fields and non-specific physical symptoms: an epidemiological study based on self-reported data and electronic medical records. *Int. J. Hyg Environ. Health* 218, 331–344.
- Baliatsas, C., van Kamp, I., Hooiveld, M., Yzermans, J., Lebre, E., 2014. Comparing non-specific physical symptoms in environmentally sensitive patients: prevalence, duration, functional status and illness behavior. *J. Psychosom. Res.* 76, 405–413.
- Baliatsas, C., Van Kamp, I., Lebre, E., Rubin, G.J., 2012. Idiopathic environmental intolerance attributed to electromagnetic fields (IEI-EMF): a systematic review of identifying criteria. *BMC Publ. Health* 12, 643.
- Bandara, P., Carpenter, D.O., 2018. Planetary electromagnetic pollution: it is time to assess its impact. *Lancet Planet. Health* 2, e512–e514.
- Bartha, L., Baumzweiger, W., Buscher, D.S., Callender, T., Dahl, K.A., Davidoff, A., Donnay, A., Edelson, S.B., Elson, B.D., Elliott, E., Flayhan, D.P., Heuser, G., Keyl, P. M., Kilburn, K.H., et al., 1999. Multiple chemical sensitivity: a 1999 consensus. *Arch. Environ. Health* 54, 147–149.
- Bas, O., Odaci, E., Kaplan, S., Acer, N., Uçok, K., Colakoglu, S., 2009. 900 MHz electromagnetic field exposure affects qualitative and quantitative features of hippocampal pyramidal cells in the adult female rat. *Brain Res.* 1265, 178–185.
- Bawin, S.M., Adey, W.R., 1976. Sensitivity of calcium binding in cerebral tissue to weak environmental electric fields oscillating at low frequency. *Proc. Natl. Acad. Sci. U. S. A.* 73, 1999–2003.
- Bell, R.D., Zlokovic, B.V., 2009. Neurovascular mechanisms and blood-brain barrier disorder in Alzheimer's disease. *Acta Neuropathol.* 118, 103–113.
- Belpomme, D., Irigaray, P., 2020. Electrohypersensitivity as a newly identified and characterized neurologic pathological disorder: how to diagnose, treat, and prevent it. *Int. J. Mol. Sci.* 21, 1915.
- Belpomme, D., Irigaray, P., 2021. Why scientifically unfounded and misleading claim should be dismissed to make true research progress in the acknowledgment of electrohypersensitivity as a new worldwide emerging pathology. *Rev. Environ. Health.* <https://doi.org/10.1515/revveh-2021-0104>.
- Belpomme, D., Campagnac, C., Irigaray, P., 2015. Reliable disease biomarkers characterizing and identifying electrohypersensitivity and multiple chemical sensitivity as two etiopathogenic aspects of a unique pathological disorder. *Rev. Environ. Health* 30, 251–271.
- Belpomme, D., Campagnac, C., Irigaray, P., 2016. Corrigendum to: reliable disease biomarkers characterizing and identifying electrohypersensitivity and multiple chemical sensitivity as two etiopathogenic aspects of a unique pathological disorder. *Rev. Environ. Health.* <https://doi.org/10.1515/revveh-2015-8888>.
- Belpomme, D., Carlo, G.L., Irigaray, P., Carpenter, D.O., Hardell, L., Kundi, M., Belyaev, I., Havas, M., Adlkofer, F., Heuser, G., Miller, A.B., Caccamo, D., De Luca, C., von Klitzing, L., Pall, M.L., Bandara, P., Stein, Y., Sage, C., Soffritti, M., Davis, D., Moskowitz, J.M., Mortazavi, S.M.J., Herbert, M.R., Moshhammer, H., Ledoit, G., Turner, R., Tweedale, A., Muñoz-Calero, P., Udasin, I., Koppel, T., Burgio, E., Vorst, A.V., 2021. The critical importance of molecular biomarkers and imaging in the study of electrohypersensitivity. A scientific consensus international report. *Int. J. Mol. Sci.* 22, 7321.
- Belpomme, D., Hardell, L., Belyaev, I., Burgio, E., Carpenter, D.O., 2018. Thermal and non-thermal health effects of low intensity non-ionizing radiation: an international perspective. *Environ. Pollut.* 242, 643–658.
- Belpomme, D., Irigaray, P., Hardell, L., Montagnier, L., Epstein, S.S., Clapp, R., Sasco, A. J., 2007. The multitude and diversity of exogenous carcinogens. *Environ. Res.* 105, 414–429.
- Belyaev, I., 2005. Non-thermal biological effects of microwaves. *Microwave Review* 11, 3–29.
- Belyaev, I.Y., Koch, C.B., Terenius, O., Roxström-Lindquist, K., Malmgren, L.O., H Sommer, W., Salford, L.G., Persson, B.R., 2006. Exposure of rat brain to 915 MHz GSM microwaves induces changes in gene expression but not double stranded DNA breaks or effects on chromatin conformation. *Bioelectromagnetics* 27, 295–306.
- Belyaev, I.Y., Kravchenko, V.G., 1994. Resonance effect of low-intensity millimeter waves on the chromatin conformational state of rat thymocytes. *Z. Naturforsch. C* 49, 352–358. <https://doi.org/10.1515/znc-1994-5-612>.
- Béres, S., Németh, A., Ajtay, Z., Kiss, I., Németh, B., Hejfel, L., 2018. Cellular phone irradiation of the head affects heart rate variability depending on inspiration/expiration ratio. *In Vivo* 32, 1145–1153.
- Bergqvist, U., Vogel, E., 1997. Possible health implications of subjective symptoms and electromagnetic fields. In: A Report Prepared by a European Group of Experts for the European Commission, DGV; Arbete Och Hälsa, vol. 19. Swedish National Institute for Working Life, Stockholm, Sweden. ISBN 91-7045-438-8.
- Bergqvist, U.O., 1984. Video display terminals and health. A technical and medical appraisal of the state of the art. *Scand. J. Work. Environ. Health* 10, 1–87.
- Blackman, C., 2009. Cell phone radiation: evidence from ELF and RF studies supporting more inclusive risk identification and assessment. *Pathophysiology* 16, 205–216.
- Blanchard, J.P., Blackman, C.F., 1994. Clarification and application of an ion parametric resonance model for magnetic field interactions with biological systems. *Bioelectromagnetics* 15, 217–238.
- Blank, M., Goodman, R., 1999. Electromagnetic fields may act directly on DNA. *J. Cell. Biochem.* 75, 369–374.
- Blank, M., Goodman, R., 2011. DNA is a fractal antenna in electromagnetic fields. *Int. J. Radiat. Biol.* 87, 409–415.

- Blank, M., 2005. Do electromagnetic fields interact with electrons in the Na,K-ATPase? *Bioelectromagnetics* 26, 677–683.
- Blettnier, M., Schlehofer, B., Breckenkamp, J., Kowall, B., Schmiedel, S., Reis, U., Potthoff, P., Schüz, J., Berg-Beckhoff, G., 2009. Mobile phone base stations and adverse health effects: phase 1 of a population-based, cross-sectional study in Germany. *Occup. Environ. Med.* 66, 118–123.
- Borgens, R.B., 1988. Stimulation of neuronal regeneration and development by steady electrical fields. In: Waxman, S.G. (Ed.), *Advances in Neurology*, 47; Functional Recovery in Neurological Disease. Raven Press, New York.
- Braune, S., Wrocklage, C., Raczek, J., Gailus, T., Lücking, C.H., 1998. Resting blood pressure increase during exposure to a radio-frequency electromagnetic field. *Lancet* 351, 1857–1858.
- Burgess, A.P., Fouquet, N.C., Seri, S., Hawken, M.B., Heard, A., Neasham, D., Little, M.P., Elliott, P., 2016. Acute Exposure to Terrestrial Trunked Radio (TETRA) has effects on the electroencephalogram and electrocardiogram, consistent with vagal nerve stimulation. *Environ. Res.* 150, 461–469.
- Cao, Z., Zhang, H., Tao, Y., Liu, J., 2000. Effects of microwave radiation on lipid peroxidation and the content of neurotransmitters in mice. *Wei Sheng Yan Jiu* 29, 28–29.
- Cardis, E., Deltour, I., Mann, S., Moissonnier, M., Taki, M., Varsier, N., Wake, K., Wiart, J., 2008. Distribution of RF energy emitted by mobile phones in anatomical structures of the brain. *Phys. Med. Biol.* 53, 2771–2783.
- Carlsson, F., Karlson, B., Ørbaek, P., Osterberg, K., Ostergren, P.O., 2005. Prevalence of annoyance attributed to electrical equipment and smells in a Swedish population, and relationship with subjective health and daily functioning. *Publ. Health* 119, 568–577.
- Carpenter, D.O., Belpomme, D., 2015. Idiopathic environmental intolerance. *Rev. Environ. Health* 30, 207.
- Carpenter, D.O., 2014. Excessive exposure to radiofrequency electromagnetic fields may cause the development of electrohypersensitivity. *Alternative Ther. Health Med.* 20, 40–42.
- Carpenter, D.O., 2015. The microwave syndrome or electro-hypersensitivity: historical background. *Rev. Environ. Health* 30, 217–222.
- Carrubba, S., Marino, A.A., 2008. The effects of low-frequency environmental-strength electromagnetic fields on brain electrical activity: a critical review of the literature. *Electromagn. Biol. Med.* 27, 83–101.
- Carrubba, S., Frilot, C., 2nd, Chesson Jr., A.L., Marino, A.A., 2010. Mobile-phone pulse triggers evoked potentials. *Neurosci. Lett.* 469, 164–168.
- Chia, S.E., Chia, H.P., Tan, J.S., 2000. Prevalence of headache among handheld cellular telephone users in Singapore: a community study. *Environ. Health Perspect.* 108, 1059–1062.
- Crasson, M., Legros, J.J., Scarpa, P., Legros, W., 1999. 50 Hz magnetic field exposure influence on human performance and psychophysiological parameters: two double-blind experimental studies. *Bioelectromagnetics* 20, 474–486.
- Croft, R.J., Chandler, J.S., Burgess, A.P., Barry, R.J., Williams, J.D., Clarke, A.R., 2002. Acute mobile phone operation affects neural function in humans. *Clin. Neurophysiol.* 113, 1623–1632.
- Croft, R.J., Hamblin, D.L., Spong, J., Wood, A.W., McKenzie, R.J., Stough, C., 2008. The effect of mobile phone electromagnetic fields on the alpha rhythm of human electroencephalogram. *Bioelectromagnetics* 29, 1–10.
- Curcio, G., Ferrara, M., Limongi, T., Tempesta, D., Di Sante, G., De Gennaro, L., Quaresima, V., Ferrari, M., 2009. Acute mobile phones exposure affects frontal cortex hemodynamics as evidenced by functional near-infrared spectroscopy. *J. Cerebr. Blood Flow Metabol.* 29, 903–910.
- Curcio, G., Ferrara, M., Moroni, F., D'Inzeo, G., Bertini, M., De Gennaro, L., 2005. Is the brain influenced by a phone call? An EEG study of resting wakefulness. *Neurosci. Res.* 53, 265–270.
- Dasdag, S., Akdag, M.Z., Erdal, M.E., Erdal, N., Ay, O.I., Ay, M.E., Yilmaz, S.G., Tasdelen, B., Yegin, K., 2015a. Long term and excessive use of 900 MHz radiofrequency radiation alter microRNA expression in brain. *Int. J. Radiat. Biol.* 91, 555–561.
- Dasdag, S., Akdag, M.Z., Erdal, M.E., Erdal, N., Ay, O.I., Ay, M.E., Yilmaz, S.G., Tasdelen, B., Yegin, K., 2015b. Effects of 2.4 GHz radiofrequency radiation emitted from Wi-Fi equipment on microRNA expression in brain tissue. *Int. J. Radiat. Biol.* 91, 555–561.
- Dasdag, S., Akdag, M.Z., Kizil, G., Kizil, M., Cakir, D.U., Yokus, B., 2012. Effect of 900 MHz radio frequency radiation on beta amyloid protein, protein carbonyl, and malondialdehyde in the brain. *Electromagn. Biol. Med.* 31, 67–74.
- Davis, S., 1997. Weak residential magnetic fields affect melatonin in humans. *Microwave News* 17 (6) novembre-décembre 1997. <http://microwavenews.com/news/backissues/n-d97issue.pdf>.
- De Iuliis, G.N., Newey, R.J., King, B.V., Aitken, R.J., 2009. Mobile phone radiation induces reactive oxygen species production and DNA damage in human spermatozoa in vitro. *PLoS One* 4, e6446.
- Del Giudice, E., Stefanini, P., Tedeschi, A., Vitiello, G., 2011. The interplay of biomolecules and water at the origin of the active behavior of living organisms. *J. Phys. Conf.* 329, 012001. Article ID: 012001.
- Deshmukh, P.S., Megha, K., Banerjee, B.D., Ahmed, R.S., Chandna, S., Abegaonkar, M.P., Tripathi, A.K., 2013. Detection of low level microwave radiation induced deoxyribonucleic acid damage vis-à-vis genotoxicity in brain of fischer rats. *Toxicol. Int.* 20, 19–24.
- Diem, E., Schwarz, C., Adlkofer, F., Jahn, O., Rüdiger, H., 2005. Non-thermal DNA breakage by mobile-phone radiation (1800 MHz) in human fibroblasts and in transformed GFSH-R17 rat granulosa cells in vitro. *Mutat. Res.* 583, 178–183.
- Dieudonné, M., 2016. Does electromagnetic hypersensitivity originate from nocebo responses? Indications from a qualitative study. *Bioelectromagnetics* 37, 14–24.
- Dodge, C., 1969. Clinical and Hygienic Aspects of Exposure to Electromagnetic Radiation. Bioscience Division of US Navy.
- Dunn, J.R., Fuller, M., Zoeger, J., Dobson, J., Heller, F., Hammann, J., Caine, E., Moskowitz, B.M., 1995. Magnetic material in the human hippocampus. *Brain Res. Bull.* 36, 149–153.
- Eberhardt, J.L., Persson, B.R., Brun, A.E., Salford, L.G., Malmgren, L.O., 2008. Blood-brain barrier permeability and nerve cell damage in rat brain 14 and 28 days after exposure to microwaves from GSM mobile phones. *Electromagn. Biol. Med.* 27, 215–229.
- Eghlidospour, M., Ghanbari, A., Mortazavi, S.M.J., Azari, H., 2017. Effects of radiofrequency exposure emitted from a GSM mobile phone on proliferation, differentiation, and apoptosis of neural stem cells. *Anat. Cell Biol.* 50, 115–123.
- Eltiti, S., Wallace, D., Russo, R., Fox, E., 2015. Aggregated data from two double blind base station provocation studies comparing individuals with idiopathic environmental intolerance with attribution to electromagnetic fields and controls. *Bioelectromagnetics* 36, 96–107.
- Eltiti, S., Wallace, D., Zougkou, K., Russo, R., Joseph, S., Rasor, P., Fox, E., 2007. Development and evaluation of the electromagnetic hypersensitivity questionnaire. *Bioelectromagnetics* 28, 137–151.
- Erickson, M.A., Banks, W.A., 2013. Blood-brain barrier dysfunction as a cause and consequence of Alzheimer's disease. *J. Cerebr. Blood Flow Metabol.* 33, 1500–1513.
- Eyvazlou, M., Zarei, E., Rahimi, A., Abazari, M., 2016. Association between overuse of mobile phones on quality of sleep and general health among occupational health and safety students. *Chronobiol. Int.* 33, 293–300.
- Forman, S.A., Holmes, C.K., McManamon, T.V., Wedding, W.R., 1982. Psychological symptoms and intermittent hypertension following acute microwave exposure. *J. Occup. Med.* 24, 932–934.
- Frei, P., Mohler, E., Braun-Fahrlander, C., Fröhlich, J., Neubauer, G., Rössli, M., QUALIFEX-team., 2012. Cohort study on the effects of everyday life radio frequency electromagnetic field exposure on non-specific symptoms and tinnitus. *Environ. Int.* 38, 29–36.
- Freude, G., Ullsperger, P., Eggert, S., Ruppe, I., 1998. Effects of microwaves emitted by cellular phones on human slow brain potentials. *Bioelectromagnetics* 19, 384–387.
- Frey, A.H., 1993. Electromagnetic field interactions with biological systems. *Faseb. J.* 7, 272–281.
- Frick, U., Rehm, J., Eichhammer, P., 2002. Risk perception, somatization, and self report of complaints related to electromagnetic fields—a randomized survey study. *Int. J. Hyg. Environ. Health* 205, 353–360.
- Funk, R.H., 2015. Endogenous electric fields as guiding cue for cell migration. *Front. Physiol.* 6, 143.
- Furtado-Filho, O.V., Borba, J.B., Maraschin, T., Souza, L.M., Henriques, J.A., Moreira, J. C., Saffi, J., 2015. Effects of chronic exposure to 950 MHz ultra-high-frequency electromagnetic radiation on reactive oxygen species metabolism in the right and left cerebral cortex of young rats of different ages. *Int. J. Radiat. Biol.* 91, 891–897.
- Furubayashi, T., Ushiyama, A., Terao, Y., Mizuno, Y., Shirasawa, K., Pongpaibool, P., Simba, A.Y., Wake, K., Nishikawa, M., Miyawaki, K., Yasuda, A., Uchiyama, M., Yamashita, H.K., Masuda, H., Hirota, S., Takahashi, M., Okano, T., Inomata-Terada, S., Sokejima, S., Maruyama, E., Watanabe, S., Taki, M., Ohkubo, C., Ugawa, Y., 2009. Effects of short-term W-CDMA mobile phone base station exposure on women with or without mobile phone related symptoms. *Bioelectromagnetics* 30, 100–113.
- Gandhi, O.P., Lazzi, G., Furse, C.M., 1996. Electromagnetic absorption in the human head and neck for mobile telephones at 835 and 1900 MHz. *IEEE Trans. Microw. Theor. Tech.* 44, 1884–1897.
- Gegeer, R.J., Foley, L.E., Casselman, A., Reppert, S.M., 2010. Animal cryptochromes mediate magnetoreception by an unconventional photochemical mechanism. *Nature* 463, 804–807.
- Genuis, S.J., 2010. Sensitivity-related illness: the escalating pandemic of allergy, food intolerance and chemical sensitivity. *Sci. Total Environ.* 408, 6047–6061.
- Ghosh, R., Yahia-Cherif, L., Hugueville, L., Ducorps, A., Lemaréchal, J.D., Thuróczy, G., de Seze, R., Selmaoui, B., 2015. Radiofrequency signal affects alpha band in resting electroencephalogram. *J. Neurophysiol.* 113, 2753–2759.
- Gibson, P.R., Kovach, S., Lupfer, A., 2015. Unmet health care needs for persons with environmental sensitivity. *J. Multidiscip. Healthc.* 8, 59–66.
- Glaser, Z.R., 1972. Bibliography of Reported Biological Phenomena ('effects') and Clinical Manifestation Attributed to Microwave and Radio-Frequency Radiation. Project MF12.524.015-00043 Report No. 2. Second Printing, with Revisions, Corrections, and Additions. Naval Medical Research Institute. National Naval Medical Center, Bethesda, Maryland 20014, U.S.A.
- Greco, F., 2020. Technical assessment of ultrasonic cerebral tomography and new scientific evaluation of its clinical interest for the diagnosis of electrohypersensitivity and multiple chemical sensitivity. *Diagnostics* 10, 427.
- Grehl, S., Martina, D., Goyenvalle, C., Deng, Z.D., Rodger, J., Sherrard, R.M., 2016. In vitro magnetic stimulation: a simple stimulation device to deliver defined low intensity electromagnetic fields. *Front. Neural Circ.* 10, 85.
- Hagström, M., Auranen, J., Ekman, R., 2013. Electromagnetic hypersensitive Finns: symptoms, perceived sources and treatments, a questionnaire study. *Pathophysiology* 20, 117–122.
- Hallberg, O., Oberfeld, G., 2006. Letter to the editor: will we all become electrosensitive? *Electromagn. Biol. Med.* 25, 189–191.
- Hardell, L., Carlberg, M., 2020. Health risks from radiofrequency radiation, including 5G, should be assessed by experts with no conflicts of interest. *Oncol. Lett.* 20, 15.
- Hardell, L., Carlberg, M., Hansson Mild, K., 2013. Use of mobile phones and cordless phones is associated with increased risk for glioma and acoustic neuroma. *Pathophysiology* 20, 85–110.

- Hardell, L., Holmberg, B., Malker, H., Paulsson, L.E., 1995. Exposure to extremely low frequency electromagnetic fields and the risk of malignant diseases—an evaluation of epidemiological and experimental findings. *Eur. J. Cancer Prev.* 4 (Suppl. 1), 3–107.
- Hardell, L., Nyberg, R., 2020. Appeals that matter or not on a moratorium on the deployment of the fifth generation, 5G, for microwave radiation. *Mol. Clin. Oncol.* 12, 247–257.
- Hardell, L., Söderqvist, F., Carlberg, M., Zetterberg, H., Mild, K.H., 2010. Exposure to wireless phone emissions and serum beta-trace protein. *Int. J. Mol. Med.* 26, 301–306.
- Havas, M., 2013. Radiation from wireless technology affects the blood, the heart, and the autonomic nervous system. *Rev. Environ. Health* 28, 75–84.
- Havas, M., Marrongelle, J., Pollner, B., Kelley, E., Rees, C., Tully, L., 2010. Provocation study using heart rate variability shows microwave radiation from 2.4 GHz cordless phone affects autonomic nervous system. Mattioli: Fidenza, Italy, 2010. In: *Non-Thermal Effects and Mechanisms of Interaction between Electromagnetic Fields and Living Matter*, pp. 273–300. ISBN 9788862611664.
- Heneka, M.T., O'Banion, M.K., 2007. Inflammatory processes in Alzheimer's disease. *J. Neuroimmunol.* 184, 69–91.
- Heuser, G., Heuser, S.A., 2017. Functional brain MRI in patients complaining of electrohypersensitivity after long term exposure to electromagnetic fields. *Rev. Environ. Health* 32, 291–299.
- Hillert, L., Berglind, N., Arnetz, B.B., Bellander, T., 2002. Prevalence of self-reported hypersensitivity to electric or magnetic fields in a population-based questionnaire survey. *Scand. J. Work. Environ. Health* 28, 33–41.
- Hocking, B., Westerman, R., 2002. Neurological changes induced by a mobile phone. *Occup. Med. (Lond.)* 52, 413–415.
- Hocking, B., Westerman, R., 2003. Neurological effects of radiofrequency radiation. *Occup. Med.* 53, 123–127.
- Hu, C., Zuo, H., Li, Y., 2021. Effects of radiofrequency electromagnetic radiation on neurotransmitters in the brain. *Front. Public Health* 9, 691880.
- Huber, R., Treyer, V., Borbély, A.A., Schuderer, J., Gottselig, J.M., Landolt, H.P., Werth, E., Berthold, T., Kuster, N., Buck, A., Achermann, P., 2002. Electromagnetic fields, such as those from mobile phones, alter regional cerebral blood flow and sleep and waking EEG. *J. Sleep Res.* 11, 289–295.
- Huber, R., Treyer, V., Schuderer, J., Berthold, T., Buck, A., Kuster, N., Landolt, H.P., Achermann, P., 2005. Exposure to pulse-modulated radio frequency electromagnetic fields affects regional cerebral blood flow. *Eur. J. Neurosci.* 21, 1000–1006.
- Huss, A., van Eijsden, M., Guxens, M., Beekhuizen, J., van Strien, R., Kromhout, H., Vrijkotte, T., Vermeulen, R., 2015. Environmental radiofrequency electromagnetic fields exposure at home, mobile and cordless phone use, and sleep problems in 7-year-old children. *PLoS One* 10, e0139869.
- Hutter, H.P., Moshhammer, H., Wallner, P., Kundl, M., 2006. Subjective symptoms, sleeping problems, and cognitive performance in subjects living near mobile phone base stations. *Occup. Environ. Med.* 63, 307–313.
- IARC (International Agency for Research on Cancer), 2002. Non-ionizing radiation, Part 1: static and extremely low-frequency (ELF) electric and magnetic fields. In: *IARC Monographs on the Evaluation of Carcinogenic Risks to Humans*, vol. 80. IARC Press, Lyon, France, p. 341.
- IARC (International Agency for Research on Cancer), 2013. Non-ionization radiation, Part 2: radiofrequency electromagnetic fields. In: *IARC Monographs on the Evaluation of Carcinogenic Risks to Humans*, vol. 102. IARC Press, Lyon, France, p. 406.
- ICNIRP (The International Commission on Non-Ionizing Radiation Protection), 1998. Guidelines for limiting exposure to time-varying electric, magnetic and electromagnetic fields (Up to 300 GHz). *Health Phys.* 74, 494–522.
- ICNIRP (The International Commission on Non-Ionizing Radiation Protection), 2010. Guidelines for limiting exposure to time-varying electric and magnetic fields (1 Hz to 100 kHz). *Health Phys.* 99, 818–836.
- ICNIRP (The International Commission on Non-Ionizing Radiation Protection), 2020. Guidelines for limiting exposure to electromagnetic fields (100 kHz to 300 GHz). *Health Phys.* 118, 483–524.
- Irigaray, P., Caccamo, D., Belpomme, D., 2018a. Oxidative stress in electrohypersensitivity self reporting patients: results of a prospective in vivo investigation with comprehensive molecular analysis. *Int. J. Mol. Med.* 42, 1885–1898.
- Irigaray, P., Lebar, P., Belpomme, D., 2018b. How ultrasonic cerebral tomography can contribute to the diagnosis of electrohypersensitivity. *J. Clin. Diagn. Res.* 6, 143.
- Ivancsits, S., Diem, E., Jahn, O., Rüdiger, H.W., 2003. Age-related effects on induction of DNA strand breaks by intermittent exposure to electromagnetic fields. *Mech. Ageing Dev.* 124, 847–850.
- Ivancsits, S., Diem, E., Pilger, A., Rüdiger, H.W., Jahn, O., 2002. Induction of DNA strand breaks by intermittent exposure to extremely-low-frequency electromagnetic fields in human diploid fibroblasts. *Mutat. Res.* 519, 1–13.
- Johansson, A., Nordin, S., Heiden, M., Sandström, M., 2010. Symptoms, personality traits, and stress in people with mobile phone-related symptoms and electromagnetic hypersensitivity. *J. Psychosom. Res.* 68, 37–45.
- Johansson, O., Gangi, S., Liang, Y., Yoshimura, K., Jing, C., Liu, P.Y., 2001. Cutaneous mast cells are altered in normal healthy volunteers sitting in front of ordinary TVs/PCs—results from open-field provocation experiments. *J. Cutan. Pathol.* 28, 513–519.
- Johnsen, S., Lohmann, K.J., 2005. The physics and neurobiology of magnetoreception. *Nat. Rev. Neurol.* 6, 703–712.
- Johnston, L., 2008. Suicides “Linked to Phone Masts”, Express.Co uk, 22 juin 2008. [www.express.co.uk/posts/view/49330/Suicides-linked-to-phone-masts](http://www.express.co.uk/posts/view/49330/Suicides-linked-to-phone-masts).
- Joubert, V., Bourthoumieu, S., Leveque, P., Yardin, C., 2008. Apoptosis is induced by radiofrequency fields through the caspase-independent mitochondrial pathway in cortical neurons. *Radiat. Res.* 169, 38–45.
- Kato, Y., Johansson, O., 2012. Reported functional impairments of electrohypersensitive Japanese: a questionnaire survey. *Pathophysiology* 19, 95–100.
- Kirschvink, J.L., Kobayashi-Kirschvink, A., Diaz-Ricci, J.C., Kirschvink, S.J., 1992b. Magnetite in human tissues: a mechanism for the biological effects of weak ELF magnetic fields. *Bioelectromagnetic. Suppl.* 1, 101–113.
- Kirschvink, J.L., Kobayashi-Kirschvink, A., Woodford, B.J., 1992a. Magnetite biomineralization in the human brain. *Proc. Natl. Acad. Sci. U.S.A.* 89, 7683–7687.
- Kleinlogel, H., Dierks, T., Koenig, T., Lehmann, H., Minder, A., Berz, R., 2008. Effects of weak mobile phone - electromagnetic fields (GSM, UMTS) on event related potentials and cognitive functions. *Bioelectromagnetics* 29, 488–497.
- Koppel, T., Vilcane, I., Ahonen, M., 2018. 50 Hz magnetic field affects heart rate variability—An experimental study, 2018. In: *Proceedings of the 2018 EMF-Med 1st World Conference on Biomedical Applications of Electromagnetic Fields (EMF-Med)*, Split, Croatia, 10–13 September 2018. IEEE, New York, NY, USA, pp. 1–2. doi.org/10.23919/EMF-MED.2018.8526072.
- Korpinen, L.H., Pääkkönen, R.J., 2009. Self-report of physical symptoms associated with using mobile phones and other electrical devices. *Bioelectromagnetics* 30, 431–437.
- Kowall, B., Breckenkamp, J., Blettner, M., Schlehofer, B., Schütz, J., Berg-Beckhoff, G., 2012. Determinants and stability over time of perception of health risks related to mobile phone base stations. *Int. J. Publ. Health* 57, 735–743.
- Krause, C.M., Sillanmäki, L., Koivisto, M., Häggqvist, A., Saarela, C., Revonsuo, A., Laine, M., Hämäläinen, H., 2000. Effects of electromagnetic field emitted by cellular phones on the EEG during a memory task. *Neuroreport* 11, 761–764.
- Lai, H., 2019. Exposure to static and extremely-low frequency electromagnetic fields and cellular free radicals. *Electromagn. Biol. Med.* 38, 231–248.
- Lai, H., 2021. Genetic effects of non-ionizing electromagnetic fields. *Electromagn. Biol. Med.* 40, 264–273.
- Lai, H., Singh, N.P., 1995. Acute low-intensity microwave exposure increases DNA single-strand breaks in rat brain cells. *Bioelectromagnetics* 16, 207–210.
- Lai, H., Singh, N.P., 2004. Magnetic-field-induced DNA strand breaks in brain cells of the rat. *Environ. Health Perspect.* 112, 687–694.
- Landgrebe, M., Frick, U., Hauser, S., Langguth, B., Rosner, R., Hajak, G., Eichhammer, P., 2008. Cognitive and neurobiological alterations in electromagnetic hypersensitive patients: results of a case-control study. *Psychol. Med.* 38, 1781–1791.
- Leone, L., Fusco, S., Mastrodonato, A., Piacentini, R., Barbat, S.A., Zaffina, S., Pani, G., Podda, M.V., Grassi, C., 2014. Epigenetic modulation of adult hippocampal neurogenesis by extremely low-frequency electromagnetic fields. *Mol. Neurobiol.* 49, 1472–1486.
- Leszczynski, D., 2021. Review of the scientific evidence on the individual sensitivity to electromagnetic fields (EHS). *Rev. Environ. Health.* <https://doi.org/10.1515/revheh-2021-0038>.
- Levallois, P., 2002. Hypersensitivity of human subjects to environmental electric and magnetic field exposure: a review of the literature. *Environ. Health Perspect.* 110, 613–618.
- Levallois, P., Neutra, R., Lee, G., Hristova, L., 2002. Study of self-reported hypersensitivity to electromagnetic fields in California. *Environ. Health Perspect.* 110, 619–623.
- Liburdy, R.P., 1992. Calcium signalling in lymphocytes and ELF fields: evidence for an electric field metric and a site of interaction involving the calcium ion channel. *FEBS Lett.* 301, 53–59.
- Lin, H., Blank, M., Goodman, R., 1999. A magnetic field-responsive domain in the human HSP70 promoter. *J. Cell. Biochem.* 75, 170–176.
- Lin, H., Blank, M., Rossol-Haseroth, K., Goodman, R., 2001. Regulating genes with electromagnetic response elements. *J. Cell. Biochem.* 81, 143–148.
- Loos, N., Thuróczy, G., Ghosn, R., Brenet-Dufour, V., Liabeuf, S., Selmaoui, B., Libert, J. P., Bach, V., Diouf, M., de Seze, R., 2013. Is the effect of mobile phone radiofrequency waves on human skin perfusion non-thermal? *Microcirculation* 20, 629–636.
- Loughran, S.P., Verrender, A., Dalecki, A., Burdon, C.A., Tagami, K., Park, J., Taylor, N. A.S., Croft, R.J., 2019. Radiofrequency electromagnetic field exposure and the resting EEG: exploring the thermal mechanism hypothesis. *Int. J. Environ. Res. Publ. Health* 16, 1505.
- Lowden, A., Akerstedt, T., Ingre, M., Wiholm, C., Hillert, L., Kuster, N., Nilsson, J.P., Arnetz, B., 2011. Sleep after mobile phone exposure in subjects with mobile phone-related symptoms. *Bioelectromagnetics* 32, 4–14.
- Lustenberger, C., Murbach, M., Dürr, R., Schmid, M.R., Kuster, N., Achermann, P., Huber, R., 2013. Stimulation of the brain with radiofrequency electromagnetic field pulses affects sleep-dependent performance improvement. *Brain Stimul.* 6, 805–811.
- Maher, B.A., Ahmed, I.A., Karloukovski, V., McLaren, D.A., Foulds, P.G., Allsop, D., Mann, D.M., Torres-Jardón, R., Calderon-Garciduenas, L., 2016. Magnetite pollution nanoparticles in the human brain. *Proc. Natl. Acad. Sci. U.S.A.* 113, 10797–10801.
- Mann, K., Röschke, J., 1996. Effects of pulsed high-frequency electromagnetic fields on human sleep. *Neuropsychobiology* 33, 41–47.
- Marková, E., Hillert, L., Malmgren, L., Persson, B.R., Belyaev, I.Y., 2005. Microwaves from GSM mobile telephones affect 53BP1 and gamma-H2AX foci in human lymphocytes from hypersensitive and healthy persons. *Environ. Health Perspect.* 113, 1172–1177.
- McCaig, C.D., Rajnicek, A.M., Song, B., Zhao, M., 2005. Controlling cell behavior electrically: current views and future potential. *Physiol. Rev.* 85, 943–978.
- McCaig, C.D., Zhao, M., 1997. Physiological electric fields modify cell behaviour. *Bioessays* 19, 819–826.

- McCarty, D.E., Carrubba, S., Chesson, A.L., Filot, C., Gonzalez-Toledo, E., Marino, A.A., 2011. Electromagnetic hypersensitivity: evidence for a novel neurological syndrome. *Int. J. Neurosci.* 121, 670–676.
- Medeiros, L.N., Sanchez, T.G., 2016. Tinnitus and cell phones: the role of electromagnetic radiofrequency radiation. *Braz. J. Otorhinolaryngol.* 82, 97–104.
- Meg Tseng, M.C., Lin, Y.P., Cheng, T.J., 2011. Prevalence and psychiatric comorbidity of self-reported electromagnetic field sensitivity in Taiwan: a population-based study. *J. Formos. Med. Assoc.* 110, 634–641.
- Megha, K., Deshmukh, P.S., Banerjee, B.D., Tripathi, A.K., Ahmed, R., Abegaonkar, M.P., 2015a. Low intensity microwave radiation induced oxidative stress, inflammatory response and DNA damage in rat brain. *Neurotoxicology* 51, 158–165.
- Megha, K., Deshmukh, P.S., Ravi, A.K., Tripathi, A.K., Abegaonkar, M.P., Banerjee, B.D., 2015b. Effect of low-intensity microwave radiation on monoamine neurotransmitters and their key regulating enzymes in rat brain. *Cell Biochem. Biophys.* 73, 93–100.
- Mild, K.H., Repacholi, M., van Deventer, E., Ravazzani, P., 2006. Electromagnetic hypersensitivity. In: *Proceedings of the WHO International Seminar and Working Group Meeting on EMF Hypersensitivity*, Prague, Czech Republic, 25–27 October 2004. World Health Organization, Geneva, Switzerland. ISBN 92-4-159412-8.
- Milde-Busch, A., von Kries, R., Thomas, S., Heinrich, S., Straube, A., Radon, K., 2010. The association between use of electronic media and prevalence of headache in adolescents: results from a population-based cross-sectional study. *BMC Neurol.* 10, 12.
- Millenbaugh, N.J., Roth, C., Sypniewska, R., Chan, V., Eggers, J.S., Kiel, J.L., Blystone, R. V., Mason, P.A., 2008. Gene expression changes in the skin of rats induced by prolonged 35 GHz millimeter-wave exposure. *Radiat. Res.* 169, 288–300.
- Miller, A.B., Sears, M.E., Morgan, L.L., Davis, D.L., Hardell, L., Oremus, M., Soskolne, C. L., 2019. Risks to health and well-being from radio-frequency radiation emitted by cell phones and other wireless devices. *Front. Public Health* 7, 223.
- Miller, C.S., 1999. Are we on the threshold of a new theory of disease? Toxicant-induced loss of tolerance and its relationship to addiction and abidction. *Toxicol. Ind. Health* 15, 284–294.
- Mohler, E., Frei, P., Fröhlich, J., Braun-Fahrlander, C., Rössli, M., QUALIFEX-team., 2012. Exposure to radiofrequency electromagnetic fields and sleep quality: a prospective cohort study. *PLoS One* 7, e37455.
- Monazzam, M.R., Hosseini, M., Matin, L.F., Aghaei, H.A., Khosroabadi, H., Hesami, A., 2014. J Sleep quality and general health status of employees exposed to extremely low frequency magnetic fields in a petrochemical complex. *Environ. Health Sci. Eng.* 12, 78.
- Mousavy, S.J., Riaz, G.H., Kamarei, M., Aliakbarian, H., Sattarahmady, N., Sharifzadeh, A., Safarian, S., Ahmad, F., Moosavi-Movahedi, A.A., 2009. Effects of mobile phone radiofrequency on the structure and function of the normal human hemoglobin. *Int. J. Biol. Macromol.* 44, 278–285.
- Muehsam, D., Lalezari, P., Lekhray, R., Abruzzo, P.M., Bolotta, A., Marini, M., Bersani, F., Aicardi, G., Pilla, A., Casper, D., 2013. Non-thermal radio frequency and static magnetic fields increase rate of hemoglobin deoxylation in a cell-free preparation. *PLoS One* 8, e61752.
- Mueller, C.H., Schierz, C., 2004. Project NEMESIS: double blind study on effects of 50 Hz EMF on sleep quality, physiological parameters and field perception in people suffering from electrical hypersensitivity. In: Mild, K.H., Repacholi, M., van Deventer, E., Ravazzani, P. (Eds.), *Electromagnetic Hypersensitivity*. Prague: Proceedings of the International Workshop on EMF Hypersensitivity, pp. 107–121.
- Navarro, E.A., Segura, J., Portoles, M., Gomez-Perretta, C., 2003. The microwave syndrome: a preliminary study in Spain. *Electromagn. Biol. Med.* 22, 161–169.
- NIEHS (National Institute of Environmental Health Sciences), 1998. Assessment of Health Effects from Exposure to Power-Line Frequency Electric and Magnetic Fields. NIEHS, Research Triangle Park, NC. Publication No. 98-3981.
- Nitby, H., Brun, A., Eberhardt, J., Malmgren, B., Persson, B.R., Salford, L.G., 2009. Increased blood-brain barrier permeability in mammalian brain 7 days after exposure to the radiation from a GSM-900 mobile phone. *Pathophysiology* 16, 103–112.
- Nordin, S., Neely, G., Olsson, D., Sandström, M., 2014. Odor and noise intolerance in persons with self-reported electromagnetic hypersensitivity. *Int. J. Environ. Res. Publ. Health* 11, 8794–8805.
- Nuccitelli, R., 1988. Ionic currents in morphogenesis. *Experientia* 44, 657–666.
- Nuccitelli, R., 2000. Endogenous electric fields during development, regeneration and wound healing. In: Costarakis, P., Stavroulakis, P. (Eds.), *Proceedings: “Millennium International Workshop on Biological Effects of Electromagnetic Fields”*. Greece, October 2000, ISBN: 960-86733-0-5.
- Oberfeld, G., Navarro, E., Portoles, M., Ceferino, M., Gomez-Perretta, C., 2004. The microwave syndrome further aspect of a Spanish study. In: *Proceedings of International Conference in Kos*, pp. 1–5 (Greece).
- Odaci, E., Hanci, H., İkinici, A., Sönmez, O.F., Aslan, A., Şahin, A., Kaya, H., Çolakoglu, S., Baş, O., 2016. Maternal exposure to a continuous 900-MHz electromagnetic field provokes neuronal loss and pathological changes in cerebellum of 32-day-old female rat offspring. *J. Chem. Neuroanat.* 75, 105–110.
- Pall, M.L., 2013. Electromagnetic fields act via activation of voltage-gated calcium channels to produce beneficial or adverse effects. *J. Cell Mol. Med.* 17, 958–965.
- Pall, M.L., 2021. Millimeter (MM) wave and microwave frequency radiation produce deeply penetrating effects: the biology and the physics. *Rev. Environ. Health.* <https://doi.org/10.1515/revheh-2020-0165>, 2021.
- Palmquist, E., Claeson, A.S., Neely, G., Stenberg, B., Nordin, S., 2014. Overlap in prevalence between various types of environmental intolerance. *Int. J. Hyg Environ. Health* 217, 427–434.
- Panagopoulos, D.J., 2017. Mobile telephony radiation effects on insect ovarian cells. The necessity for real exposures bioactivity assessment. The key role of polarization, and the “Ion Forced-Oscillation Mechanism”. In: Geddes, C.D. (Ed.), *Microwave Effects on DNA and Proteins*. Springer.
- Panagopoulos, D.J., 2019. Comparing DNA damage induced by mobile telephony and other types of man-made electromagnetic fields. *Mutat. Res. Rev. Mutat. Res.* 781, 53–62.
- Panagopoulos, D.J., Chavdoula, E.D., Nezis, I.P., Margaritis, L.H., 2007. Cell death induced by GSM 900MHz and DCS 1800MHz mobile telephony radiation. *Mutat. Res.* 626, 69–78.
- Panagopoulos, D.J., Johansson, O., Carlo, G.L., 2015a. Polarization: a key difference between man-made and natural electromagnetic fields, in regard to biological activity. *Sci. Rep.* 12, 14914.
- Panagopoulos, D.J., Johansson, O., Carlo, G.L., 2015b. Real versus simulated mobile phone exposures in experimental studies. *BioMed Res. Int.* 607053.
- Panagopoulos, D.J., Karabarbounis, A., Margaritis, L.H., 2002. Mechanism for action of electromagnetic fields on cells. *Biochem. Biophys. Res. Commun.* 298, 95–102.
- Panagopoulos, D.J., Karabarbounis, A., Yakymenko, I., Chrousos, G.P., 2021. Human-made electromagnetic fields: ion forced-oscillation and voltage-gated ion channel dysfunction, oxidative stress and DNA damage (Review). *Int. J. Oncol.* 59, 92. <https://doi.org/10.3892/ijo.2021.5272>.
- Patel, J.P., Frey, B.N., 2015. Radiofrequency electromagnetic field exposure and non-specific symptoms of ill health: a systematic review. *Disruption in the Blood-Brain Barrier: The Missing Link between Brain and Body Inflammation in Rössli, M., (2008). Environ. Res.* 107, 277–287.
- Perry, F.S., Reichmanis, M., Marino, A.A., Becker, R.O., 1981. Environmental power-frequency magnetic fields and suicide. *Health Phys.* 41, 267–277.
- Phillips, J.L., Singh, N.P., Lai, H., 2009. Electromagnetic fields and DNA damage. *Pathophysiology* 16, 79–88.
- Pollack, H., 1979. The microwave syndrome. *Bull. N. Y. Acad. Med.* 55, 1240–1243.
- Poole, C., Kavet, R., Funch, D.P., Donelan, K., Charry, J.M., Dreyer, N.A., 1993. Depressive symptoms and headaches in relation to proximity of residence to an alternating-current transmission line right-of-way. *Am. J. Epidemiol.* 137, 318–330.
- Randolph, T.G., 1962. *Human Ecology and Susceptibility to the Chemical Environment*. Charles C Thomas, Springfield, IL, p. 148pp.
- Rea, W.J., Pan, Y., Fenyves, E.F., Sujişawa, I., Suyama, H., Samadi, N., Ross, G.H., 1991. Electromagnetic field sensitivity. *J. Bioelectr.* 10, 214–256.
- Redmayne, M., Johansson, O., 2014. Could myelin damage from radiofrequency electromagnetic field exposure help explain the functional impairment electrohypersensitivity? A review of the evidence. *J. Toxicol. Environ. Health B Crit. Rev.* 17, 247–258.
- Reeves, G.L., 2000. Review of extensive workups of 34 patients over-exposed to radiofrequency radiation. *Aviat Space Environ. Med.* 71, 206–215.
- Report of the Workshop on Multiple Chemical Sensitivities (MCS), Berlin, Germany, (21–23 February 1996): <https://apps.who.int/iris/handle/10665/26723/browse?authority=Multiple+Chemical+Sensitivity&type=mesh>.
- Roggeveen, S., van Os, J., Viechtbauer, W., Lousberg, R., 2015. EEG changes due to experimentally induced 3G mobile phone radiation. *PLoS One* 10, e0129496.
- Rössli, M., 2008. Radiofrequency electromagnetic field exposure and non-specific symptoms of ill health: a systematic review. *Environ. Res.* 107, 277–287.
- Rössli, M., Frei, P., Mohler, E., Hug, K., 2010a. Systematic review on the health effects of exposure to radiofrequency electromagnetic fields from mobile phone base stations. *Bull. World Health Organ.* 88, 887–896.
- Rössli, M., Mohler, E., Frei, P., 2010b. Sense and sensibility in the context of radiofrequency electromagnetic field exposure. *C.R. physique.* 11, 576–584.
- Rubin, G.J., Nieto-Hernandez, R., Wessely, S., 2010. Idiopathic environmental intolerance attributed to electromagnetic fields (formerly ‘electromagnetic hypersensitivity’): an updated systematic review of provocation studies. *Bioelectromagnetics* 31, 1–11.
- Rubin, G.J., Hillert, L., Nieto-Hernandez, R., van Rongen, E., Oftedal, G., 2011. Do people with idiopathic environmental intolerance attributed to electromagnetic fields display physiological effects when exposed to electromagnetic fields? A systematic review of provocation studies. *Bioelectromagnetics* 32, 593–609.
- Saili, L., Hanini, A., Smirani, C., Azzouz, I., Azzouz, A., Sakly, M., Abdelmelek, H., Bouslama, Z., 2015. Effects of acute exposure to WIFI signals (2.45GHz) on heart variability and blood pressure in Albinos rabbit. *Environ. Toxicol. Pharmacol.* 40, 600–605.
- Salford, L.G., Brun, A.E., Eberhardt, J.L., Marmgren, L., Persson, B.R., 2003. Nerve cell damage in mammalian brain after exposure to microwaves from GSM mobile phones. *Environ. Health Perspect.* 111, 881–883.
- Salford, L.G., Brun, A., Sturesson, K., Eberhardt, J.L., Persson, B.R., 1994. Permeability of the blood-brain barrier induced by 915 MHz electromagnetic radiation, continuous wave and modulated at 8, 16, 50, and 200 Hz. *Microsc. Res. Tech.* 27, 535–542.
- Santini, R., Santini, P., LeRuz, P., Danze, J.M., Seigne, M., 2003. Survey study of people living in the vicinity of cellular phone base stations. *Electromagn. Biol. Med.* 22, 41–49.
- Santini, R., Seigne, M., Bonhomme-Faivre, L., Bouet, S., Defrasme, E., Sage, M., 2002. Symptoms experienced by users of digital cellular phones: a study of a French engineering school. *Electromagn. Biol. Med.* 21, 81–88.
- Schilling, C.J., 1997. Effects of acute exposure to ultrahigh radiofrequency radiation on three antenna engineers. *Occup. Environ. Med.* 54, 281–284.
- Schilling, C.J., 2000. Effects of exposure to very high frequency radiofrequency radiation on six antenna engineers in two separate incidents. *Occup. Med.* 50, 49–56.
- Schmid, M.R., Loughran, S.P., Regel, S.J., Murbach, M., Bratic Grunauer, A., Rusterholz, T., Bersagliere, A., Kuster, N., Achermann, P., 2012. Sleep EEG alterations: Effects of different pulse-modulated radio frequency electromagnetic fields. *J. Sleep Res.* 21, 50–58.

- Schmiedchen, K., Driessen, S., Oftedal, G., 2019. Methodological limitations in experimental studies on symptom development in individuals with idiopathic environmental intolerance attributed to electromagnetic fields (IEI-EMF) - a systematic review. *Environ. Health*. 18, 88.
- Schreier, N., Huss, A., Rössli, M., 2006. The prevalence of symptoms attributed to electromagnetic field exposure: a cross-sectional representative survey in Switzerland. *Sozial- Präventivmed*. 51, 202–209.
- Schröttner, J., Leitgeb, N., 2008. Sensitivity to electricity—temporal changes in Austria. *BMC Publ. Health* 8, 310.
- Schüz, J., Petters, C., Egle, U.T., Jansen, B., Kimbel, R., Letzel, S., Nix, W., Schmidt, L.G., Vollrath, L., 2006. The "Mainzer EMF-Wachhund": results from a watchdog project on self-reported health complaints attributed to exposure to electromagnetic fields. *Bioelectromagnetics* 27, 280–287.
- Sekeroglu, V., Akar, A., Sekeroglu, Z.A., 2012. Cytotoxic and genotoxic effects of highfrequency electromagnetic fields (GSM 1800 MHz) on immature and mature rats. *Ecotoxicol. Environ. Saf.* 80, 140–144.
- Sekeroglu, Z.A., Akar, A., Sekeroglu, V., 2013. Evaluation of the cytogenetic damage in immature and mature rats exposed to 900 MHz radiofrequency electromagnetic fields. *Int. J. Radiat. Biol.* 89, 985, 892.
- Shi, R., Borgens, R.B., 1995. Three-dimensional gradients of voltage during development of the nervous system as invisible coordinates for the establishment of embryonic pattern. *Dev. Dynam.* 202, 101–114.
- Silva, D.F., Barros, W.R., Almeida Mda, C., Régo, M.A., 2015. Exposure to non-ionizing electromagnetic radiation from mobile telephony and the association with psychiatric symptoms. *Cad0 Saude Publica*. 31, 2110–2126.
- Sirav, B., Seyhan, N., 2016. Effects of GSM modulated radio-frequency electromagnetic radiation on permeability of blood-brain barrier in male & female rats. *J. Chem. Neuroanat.* 75, 123–127.
- Sonmez, O.F., Odaci, E., Bas, O., Kaplan, S., 2010. Purkinje cell number decreases in the adult female rat cerebellum following exposure to 900 MHz electromagnetic field. *Brain Res.* 1356, 95–101.
- Sosa, M., Bernal-Alvarado, J., Jiménez-Moreno, M., Hernández, J.C., Gutiérrez-Juárez, G., Vargas-Luna, M., Huerta, R., Villagómez-Castro, J.C., Palomares, P., 2005. Magnetic field influence on electrical properties of human blood measured by impedance spectroscopy. *Bioelectromagnetics* 26, 564–570.
- Stein, Y., Udasin, I.G., 2020. Electromagnetic hypersensitivity (EHS, microwave syndrome) - review of mechanisms. *Environ. Res.* 186, 109445.
- Sudan, M., Kheifets, L., Arah, O., Olsen, J., Zeltzer, L., 2012. Prenatal and postnatal cell phone exposures and headaches in children. *Open Pediatr. Med. J.* 6, 46–52.
- Tan, S.Z., Tan, P.C., Luo, L.Q., Yang, Z.L., Zhao, X.L., Zhao, L., Dong, J., Zhang, J., Yao, B. W., Xu, X.P., Tian, G., Chen, J.K., Wang, H., Peng, R.Y., 2019. Exposure effects of terahertz waves on primary neurons and neuron-like cells under nonthermal conditions. *Biomed. Environ. Sci.* 32, 739–754.
- Tao, R., Huang, K., 2011. Reducing blood viscosity with magnetic fields. *Phys. Rev. E. Stat. Nonlin. Soft. Matter. Phys.* 84, 011905.
- Teimori, F., Khaki, A.A., Rajabzadeh, A., Roshangar, L., 2016. The effects of 30 mT electromagnetic fields on hippocampus cells of rats. *Surg. Neurol. Int.* 7, 70.
- Tettamanti, G., Auvinen, A., Åkerstedt, T., Kojo, K., Ahlbom, A., Heinävaara, S., Elliott, P., Schüz, J., Deltour, I., Kromhout, H., Toledano, M.B., Poulsen, A.H., Johansen, C., Vermeulen, R., Feychting, M., Hillert, L., COSMOS Study Group, 2020. Long-term effect of mobile phone use on sleep quality: results from the cohort study of mobile phone use and health (COSMOS). *Environ. Int.* 140, 105687.
- Trimmel, M., Schweiger, E., 1998. Effects of an ELF (50 Hz, 1 mT) electromagnetic field (EMF) on concentration in visual attention, perception and memory including effects of EMF sensitivity. *Toxicol. Lett.* 96–97, 377–382.
- Tuengler, A., von Klitzing, L., 2013. Hypothesis on how to measure electromagnetic hypersensitivity. *Electromagn. Biol. Med.* 32, 281–290.
- Valentini, E., Ferrara, M., Presaghi, F., De Gennaro, L., Curcio, G., 2010. Systematic review and meta-analysis of psychomotor effects of mobile phone electromagnetic fields. *Occup. Environ. Med.* 67, 708–716.
- van Dongen, D., Smid, T., Timmermans, D.R., 2014. Symptom attribution and risk perception in individuals with idiopathic environmental intolerance to electromagnetic fields and in the general population. *Perspect. Public Health.* 134, 160–168.
- Vander Vorst, A., Rosen, A., Kotsuka, Y., 2006. RF-microwave Interaction with Biological Tissues. Wiley IEE Press, 2006.
- Vecchio, F., Buffo, P., Sergio, S., Iacoviello, D., Rossini, P.M., Babiloni, C., 2012. Mobile phone emission modulates event-related desynchronization of  $\alpha$  rhythms and cognitive-motor performance in healthy humans. *Clin. Neurophysiol.* 123, 121–128.
- Verkasalo, P.K., Kaprio, J., Varjonen, J., Romanov, K., Heikkilä, K., Koskenvuo, M., 1997. Magnetic fields of transmission lines and depression. *Am. J. Epidemiol.* 146, 1037–1045.
- Verrender, A., Loughran, S.P., Dalecki, A., McKenzie, R., Croft, R.J., 2016. Pulse modulated radiofrequency exposure influences cognitive performance. *Int. J. Radiat. Biol.* 92, 603–610.
- Volkow, N.D., Tomasi, D., Wang, G.J., Vaska, P., Fowler, J.S., Telang, F., Alexoff, D., Logan, J., Wong, C., 2011. Effects of cell phone radiofrequency signal exposure on brain glucose metabolism. *JAMA* 305, 808–813.
- von Klitzing, L., 1995. Low-Frequency pulsed electromagnetic fields influence EEG of man. *Phys. Med. XI*, 77–80.
- von Klitzing, L., 2021. Artificial EMG by WLAN-exposure. *J. Biostat. Biometric. App.* 6 (101). ISSN 2455-765X.
- Wagner, P., Rösche, J., Mann, K., Fell, J., Hiller, W., Frank, C., Grözing, M., 2000. Human sleep EEG under the influence of pulsed radio frequency electromagnetic fields. Results from polysomnographies using submaximal high power flux densities. *Neuropsychobiology* 42, 207–412.
- Walleczek, J., 1992. Electromagnetic field effects on cells of the immune system: the role of calcium signaling. *Faseb. J.* 6, 3177–3185.
- Wang, C.X., Hilburn, I.A., Wu, D.A., Mizuhara, Y., Cousté, C.P., Abrahams, J.N.H., Bernstein, S.E., Matani, A., Shimojo, S., Kirschvink, J.L., 2019. Transduction of the geomagnetic field as evidenced from alpha-band Activity in the human brain. *eNeuro* 26. ENEURO.0483-18.2019.
- Weisenseel, M.H., 1983. Control of differentiation and growth by endogenous electric currents. In: Hoppe, W., Lohmann, W., Markl, H., Ziegler, H. (Eds.), *Biophysics*. Springer –Verlag, Berlin, pp. 460–465.
- WHO (World Health Organization), 2005. Electromagnetic Fields and Public Health, Electromagnetic Hypersensitivity; WHO Fact Sheet No. 296. World Health Organization, Geneva, Switzerland.
- WHO (World Health Organization), 2006. Framework for Developing Health-Based EMF Standards. WHO, Geneva, Switzerland, 2006; ISBN 9241594330. [www.who.int/pe-h-emf/standards/EMF\\_standards\\_framework%5b1%5d.pdf](http://www.who.int/pe-h-emf/standards/EMF_standards_framework%5b1%5d.pdf).
- WHO (World Health Organization), 2014. Electromagnetic Fields and Public Health: Mobile Phones; Fact Sheet No.193. World Health Organization, Geneva, Switzerland.
- Williams, R.A., Webb, T.S., 1980. Exposure to radiofrequency radiation from an aircraft radar unit. *Aviat Space Environ. Med.* 51, 1243–1244.
- Yakymenko, I., Tsybulin, O., Sidorik, E., Henshel, D., Kyrylenko, O., Kyrylenko, S., 2016. Oxidative mechanisms of biological activity of low-intensity radiofrequency radiation. *Electromagn. Biol. Med.* 35, 186–202.
- Yang, X.S., He, G.L., Hao, Y.T., Xiao, Y., Chen, C.H., Zhang, G.B., Yu, Z.P., 2012. Exposure to 2.45 GHz electromagnetic fields elicits an HSP-related stress response in rat hippocampus. *Brain Res. Bull.* 88, 371–378.
- Yao, L., McCaig, C.D., Zhao, M., 2009. Electrical signals polarize neuronal organelles, direct neuron migration, and orient cell division. *Hippocampus* 19, 855–868.
- Zaret, M.M., 1973. Microwave cataracts. *Med. Trial Tech. Q.* 19, 246. Z52.
- Zhang, Y., Li, Z., Gao, Y., Zhang, C., et al., 2015. Effects of fetal microwave radiation exposure on offspring behavior in mice. *J. Radiat. Res.* 56, 261–268. <https://doi.org/10.1093/jrr/rru097>.
- Zheng, F., Gao, P., He, M., Li, M., Tan, J., Chen, D., Zhou, Z., Yu, Z., Zhang, L., 2015. Association between mobile phone use and self-reported well-being in children: a questionnaire-based cross-sectional study in Chongqing, China. *BMJ Open* 5, e007302.
- Zuo, H., Lin, T., Wang, D., Peng, R., Wang, S., Gao, Y., Xu, X., Li, Y., Wang, S., Zhao, L., Wang, L., Zhou, H., 2014. Neural cell apoptosis induced by microwave exposure through mitochondria-dependent caspase-3 pathway. *Int. J. Med. Sci.* 11, 426–435.