

BODY SCAN OF MILLICENT BLACK by Melinda Kidder, PI 1/19/2014




SCAN OBSERVATIONS OF NOTE

Using Method Standard One, with Black lying prone, the ELF/EMF Scan readings varied. The standardfor the room was 0.00-0.01µ. Testing on a brewing coffee pot in an adjacent room showed 0.32 µ.

With equipment showing readings within normal limits, testing proceeded. Normal for Black was 0.00-0.01µ except at the following locations: 0.02µ at crown of the skull; 0.02µ at the Ⓛ parietal bone (forehead); 0.02µ at the center of the parietal (forehead); 0.02µ at the Ⓡ trapezious; 0.02µ-0.05µ oscillating at the Ⓡ pectoralis minor; 0.02µ at the Ⓡ deltoid; 0.02µ at the Ⓡ medial bicep; 0.02µ at six points in what appears to be a line starting from mid sternum, past a scar, and ending at the pubic bone; 0.02µ-0.05µ oscillating at the Ⓛ lower quadrant of the abdomen; 0.02µ in three points that appear to be a line across the Ⓛ hip flexor; 0.02µ at the Ⓛ wrist; 0.02µ at the Ⓛ rectus femorus (quadriceps); 0.02µ at the Ⓛ knee with two points superior and one point inferior; 0.02µ with one point superior and one point inferior to the Ⓡ knee; 0.02µ-0.04µ oscillating just inferior to the Ⓡ patella; 0.02µ-0.05µ oscillating at the Ⓡ distal phalanges of the foot; 0.02µ at the median of the Ⓛ tibia; 0.02µ at the fifth distal phalange of the Ⓛ foot; -0.00-0.02µ oscillating at the shoulders, bilaterally (negative readings are unusual in this investigator's experience); 0.05µ at the Ⓛ scapula; 0.02µ at the Ⓛ tricep; 0.03µ at the Ⓛ brachioradialis (forearm muscle); 0.03µ at the Ⓡ kidney; 0.02µ at the Ⓡ brachioradialis (forearm muscle); 0.02µ, bilaterally, at the TrP2 of the gluteus maximus; 0.02µ-0.04µ oscillating at the Ⓡ gracillis (groin); 0.02µ at the Ⓡ lateral cutaneous nerve of the thigh; 0.02µ at the Ⓛ median of the semitendinosis (hamstring); 0.02µ-0.08µ oscillating, at the Ⓛ inferior semitendinosis; 0.02µ in a band across the median of the Ⓡ semitendinosis; 0.02µ in a band across the inferior of the Ⓡ semitendinosis; 0.02µ at the Ⓛ biceps femoris, short head; 0.02µ at the Ⓛ inferior semitendinosis; 0.02µ at the Ⓡ inferior semitendinosis; 0.02µ at the Ⓡ lateral cutaneous nerve of the calf; 0.02µ at the Ⓡ Achilles tendon; 0.02µ at the Ⓡ cuboid bone of the foot; 0.03µ at the Ⓛ cuboid bone of the foot. An image reflecting these points is shown on the following page:





SCAN OBSERVATIONS OF NOTE

Using Method Standard One, with Black lying prone, the ELF/EMF Scan readings varied. The standardfor the room was 0.00-0.01µ. Testing on a brewing coffee pot in an adjacent room showed 0.32 µ.

With equipment showing readings within normal limits, testing proceeded. Normal for Black was 0.00-0.01µ except at the following locations: 0.02µ at crown of the skull; 0.02µ at the Ⓛ parietal bone (forehead); 0.02µ at the center of the parietal (forehead); 0.02µ at the ® trapezious; 0.02µ-0.05µ oscillating at the ® pectoralis minor; 0.02µ at the ® deltoid; 0.02µ at the ® medial bicep; 0.02µ at six points in what appears to be a line starting from mid sternum, past a scar, and ending at the pubic bone; 0.02µ-0.05µ oscillating at the Ⓛ lower quadrant of the abdomen; 0.02µ in three points that appear to be a line across the Ⓛ hip flexor; 0.02µ at the Ⓛ wrist; 0.02µ at the Ⓛ rectus femorus (quadriceps); 0.02µ at the Ⓛ knee with two points superior and one point inferior; 0.02µ with one point superior and one point inferior to the ® knee; 0.02µ-0.04µ oscillating just inferior to the ® patella; 0.02µ-0.05µ oscillating at the ® distal phalanges of the foot; 0.02µ at the median of the Ⓛ tibia; 0.02µ at the fifth distal phalange of the Ⓛ foot; -0.00-0.02µ oscillating at the shoulders, bilaterally (negative readings are unusual in this investigator's experience); 0.05µ at the Ⓛ scapula; 0.02µ at the Ⓛ tricep; 0.03µ at the Ⓛ brachioradialis (forearm muscle); 0.03µ at the ® kidney; 0.02µ at the ® brachioradialis (forearm muscle); 0.02µ, bilaterally, at the TrP2 of the gluteus maximus; 0.02µ-0.04µ oscillating at the ® gracillis (groin); 0.02µ at the ® lateral cutaneous nerve of the thigh; 0.02µ at the Ⓛ median of the semitendinosis (hamstring); 0.02µ-0.08µ oscillating, at the Ⓛ inferior semitendinosis; 0.02µ in a band across the median of the ® semitendinosis; 0.02µ in a band across the inferior of the ® semitendinosis; 0.02µ at the Ⓛ biceps femoris, short head; 0.02µ at the Ⓛ inferior semitendinosis; 0.02µ at the ® inferior semitendinosis; 0.02µ at the ® lateral cutaneous nerve of the calf; 0.02µ at the ® Achilles tendon; 0.02µ at the ® cuboid bone of the foot; 0.03µ at the Ⓛ cuboid bone of the foot. An image reflecting these points is shown on the following page: