APPENDIX B — SCIENTIFIC REFERENCES OF SYNERGISTIC EFFECTS

Belyaev, I. Y., Bauréus Koch, C., Terenius, O., Roxström-Lindquist, K., Malmgren, L. O. G., Sommer, W. H., Salford, L. G., & Persson, B. R. R. (2006). Exposure of rat brain to 915 MHz GSM microwaves induces changes in gene expression but not double stranded DNA breaks or effects on chromatin conformation. *Bioelectromagnetics*, 27(4), 295–306. https://doi.org/10.1002/bem.20216

Byun, Y.-H., Ha, M., Kwon, H.-J., Hong, Y.-C., Leem, J.-H., Sakong, J., Kim, S. Y., Lee, C. G., Kang, D., Choi, H.-D., & Kim, N. (2013). Mobile phone use, blood lead levels, and attention deficit hyperactivity symptoms in children: A longitudinal study. *PLoS ONE*, 8(3), e59742. https://doi.org/10.1371/journal.pone.0059742

Eberhardt, J. L., Persson, B. R. R., Brun, A. E., Salford, L. G., & Malmgren, L. O. G. (2008). Blood-brain barrier permeability and nerve cell damage in rat brain 14 and 28 days after exposure to microwaves from GSM mobile phones. *Electromagnetic Biology and Medicine*, 27(3), 215–229. https://doi.org/10.1080/15368370802344037

International Agency for Research on Cancer. (2013). Non-ionizing radiation, Part 2: Radiofrequency electromagnetic fields (IARC Monographs on the Evaluation of Carcinogenic Risks to Humans, Vol. 102). World Health Organization.

Juutilainen, J., Kumlin, T., & Naarala, J. (2006). Do extremely low frequency magnetic fields enhance the effects of environmental carcinogens? A meta-analysis of experimental studies. *International Journal of Radiation Biology*, 82(1), 1–12. https://doi.org/10.1080/09553000600577839

Kostoff, R. N., & Lau, C. G. Y. (2013). Combined biological and health effects of electromagnetic fields and other agents in the published literature. *Technological Forecasting and Social Change*, 80(7), 1331–1349. https://doi.org/10.1016/j.techfore.2012.12.006

Kostoff, R. N., & Lau, C. G. Y. (2017). Modified health effects of non-ionizing electromagnetic radiation combined with other agents reported in the biomedical literature. *In Microwave effects on DNA and proteins* (pp. 97–158).

Lai, H. (1997, September 16–17). Neurological effects of radiofrequency electromagnetic radiation relating to wireless communication technology [Conference presentation]. IBC-UK Conference: Mobile Phones—Is There a Health Risk?

Lai, H. (2021). Genetic effects of non-ionizing electromagnetic fields. *Electromagnetic Biology and Medicine*, 40(2), 264–273. https://doi.org/10.1080/15368378.2021.1881866

Lerchl, A., Klose, M., Grote, K., Wilhelm, A. F. X., Spathmann, O., Fiedler, T., Streckert, J., Hansen, V., & Clemens, M. (2015). Tumor promotion by exposure to radiofrequency electromagnetic fields below exposure limits for humans. *Biochemical and Biophysical Research Communications*, 459(4), 585–590. https://doi.org/10.1016/j.bbrc.2015.02.151

Leszczynski, D., Joenväärä, S., Reivinen, J., & Kuokka, R. (2002). Non-thermal activation of the hsp27/p38MAPK stress pathway by mobile phone radiation in human endothelial cells: Molecular mechanism for cancer- and blood-brain barrier-related effects. *Differentiation*, 70(2–3), 120–129. https://doi.org/10.1046/j.1432-0436.2002.700207.x

López-Álvarez, C., López-Martín, M. E., Rodríguez-González, J. A., & Ares-Pena, F. J. (2023). A review on the effects of thermal inversions and electromagnetic fields on cell cultures and wireless communications. *Sensors*, 23(23), 9567. https://doi.org/10.3390/s23239567

Mariea, T. J., & Carlo, G. L. (2007). Wireless radiation in the etiology and treatment of autism: Clinical observations and mechanisms. *Journal of the Australasian College of Nutritional and Environmental Medicine*, 26(2), 3–7. https://galweiss.com/wp-content/uploads/2021/08/Wireless-Radiation-in-the-Etiology-and-treatment-of-autism-Mariea-and-Carlo.pdf

McCredden, J., McLean, L., & Steinemann, A. (2026). Wireless sensitivity and co-morbidities: A prevalence study in Australia, Canada, and the United States. *Next Research, 8*, Article 101577. https://doi.org/10.1016/j.nexres.2026.101577

Nittby, H., Brun, A., Eberhardt, J., Malmgren, L., Persson, B. R. R., & Salford, L. G. (2009). Increased blood-brain barrier permeability in mammalian brain 7 days after exposure to the radiation from a GSM-900 mobile phone. *Pathophysiology*, 16(2–3), 103–112. https://doi.org/10.1016/j.pathophys.2009.01.001

Nittby, H., Grafström, G., Eberhardt, J. L., Malmgren, L., Brun, A., Persson, B. R. R., & Salford, L. G. (2008). Radiofrequency and extremely low-frequency electromagnetic field effects on the blood-brain barrier. *Electromagnetic Biology and Medicine*, 27(2), 103–126. https://doi.org/10.1080/15368370802061995

Omar, A. I. (2014). Tumor treating field therapy in combination with bevacizumab for the treatment of recurrent glioblastoma. *Journal of Visualized Experiments*, 92, e51638. https://doi.org/10.3791/51638

Persson, B. R. R., Salford, L. G., Brun, A., Eberhardt, J. L., & Malmgren, L. (1992). Increased permeability of the blood-brain barrier induced by magnetic and electromagnetic fields. *Annals of the New York Academy of Sciences*, 649, 356–358. https://doi.org/10.1111/j.1749-6632.1992.tb49629.x

Poulletier de Gannes, F., Masuda, H., Billaudel, B., Poque-Haro, E., Hurtier, A., Lévêque, P., Ruffié, G., Taxile, M., Veyret, B., & Lagroye, I. (2017). Effects of GSM and UMTS mobile telephony signals on neuron degeneration and blood-brain barrier permeation in the rat brain. *Scientific Reports*, 7, 15496. https://doi.org/10.1038/s41598-017-15690-1

Rajkovic, V., Matavulj, M., & Johansson, O. (2010). Combined exposure of peripubertal male rats to the endocrine-disrupting compound atrazine and power-frequency electromagnetic fields causes degranulation of cutaneous mast cells: A new toxic environmental hazard? *Archives of Environmental Contamination and Toxicology*, 59(2), 334–341. https://doi.org/10.1007/s00244-010-9477-6

Ruiz-Gómez, M. J., & Martínez-Morillo, M. (2005). Enhancement of the cell-killing effect of ultraviolet-C radiation by short-term exposure to a pulsed magnetic field. *International Journal of Radiation Biology*, 81(7), 483–490. https://doi.org/10.1080/09553000500196805

Salford, L. G., Brun, A., Eberhardt, J. L., Malmgren, L., & Persson, B. R. R. (2003). Nerve cell damage in mammalian brain after exposure to microwaves from GSM mobile phones. *Environmental Health Perspectives*, 111(7), 881–883. https://doi.org/10.1289/ehp.6039

Salford, L. G., Brun, A., Sturesson, K., Eberhardt, J. L., & Persson, B. R. R. (1994). Permeability of the blood-brain barrier induced by 915 MHz electromagnetic radiation, continuous wave and modulated at 8, 16, 50, and 200 Hz. Microscopy Research and Technique, 27(6), 535–542. https://doi.org/10.1002/jemt.1070270608

Soffritti, M., & Giuliani, L. (2019). The carcinogenic potential of non-ionizing radiations: The cases of S-50 Hz MF and 1.8 GHz GSM radiofrequency radiation. *Basic & Clinical Pharmacology & Toxicology*, 125(Suppl. 3), 58–69. https://doi.org/10.1111/bcpt.13215

Soffritti, M., Tibaldi, E., Padovani, M., Hoel, D. G., Giuliani, L., Bua, L., Lauriola, M., Falcioni, L., Manservigi, M., Manservisi, F., Panzacchi, S., & Belpoggi, F. (2016). Life-span exposure to sinusoidal-50 Hz magnetic field and acute low-dose γ radiation induce carcinogenic effects in Sprague-Dawley rats. *International Journal of Radiation Biology*, 92(4), 202–214. https://doi.org/10.3109/09553002.2016.1144942

Soffritti, M., Tibaldi, E., Padovani, M., Hoel, D. G., Giuliani, L., Bua, L., Lauriola, M., Falcioni, L., Manservigi, M., Manservisi, F., & Belpoggi, F. (2016). Synergism between sinusoidal-50 Hz magnetic field and formaldehyde in triggering carcinogenic effects in male Sprague-Dawley rats. *American Journal of Industrial Medicine*, 59(7), 509–521. https://doi.org/10.1002/ajim.22598

Sueiro-Benavides, R. A., Leiro-Vidal, J. M., Salas-Sánchez, Á. Á., Rodríguez-González, J. A., & Ares-Pena, F. J. (2021). Radiofrequency at 2.45 GHz increases toxicity, pro-inflammatory and pre-apoptotic activity caused by black carbon in the RAW 264.7 macrophage cell line. *Science of the Total Environment*, 765, 142681. https://doi.org/10.1016/j.scitotenv.2020.142681

Szmigielski, S., Szudzinski, A., Pietraszek, A., Bielec, M., Janiak, M., & Wrembel, J. K. (1982). Accelerated development of spontaneous and benzopyrene-induced skin cancer in mice exposed to 2450-MHz microwave radiation. *Bioelectromagnetics*, 3(2), 179–191. https://doi.org/10.1002/bem.2250030202

Szudziński, A., Pietraszek, A., Janiak, M., Wrembel, J., Kałczak, M., & Szmigielski, S. (1982). Acceleration of the development of benzopyrene-induced skin cancer in mice by microwave radiation. *Archives of Dermatological Research*, 274(3–4), 303–312. https://doi.org/10.1007/BF00403734

Tang, J., Zhang, Y., Yang, L., Chen, Q., Tan, L., Zuo, S., Feng, H., Chen, Z., & Zhu, G. (2015). Exposure to 900 MHz electromagnetic fields activates the mkp-1/ERK pathway and causes blood-brain barrier damage and cognitive impairment in rats. *Brain Research*, 1601, 92–101. https://doi.org/10.1016/j.brainres.2015.01.019

Tillmann, T., Ernst, H., Ebert, S., Kuster, N., Behnke, W., Rittinghausen, S., & Dasenbrock, C. (2007). Carcinogenicity study of GSM and DCS wireless communication signals in B6C3F1 mice. *Bioelectromagnetics*, 28(3), 173–187. https://doi.org/10.1002/bem.20283

Tillmann, T., Ernst, H., Streckert, J., Zhou, Y., Taugner, F., Hansen, V., & Dasenbrock, C. (2010). Indication of cocarcinogenic potential of chronic UMTS-modulated radiofrequency exposure in an ethylnitrosourea mouse model. *International Journal of Radiation Biology*, 86(7), 529–541. https://doi.org/10.3109/09553001003734501

Turner, M. C., Benke, G., Bowman, J. D., Figuerola, J., Fleming, S., Hours, M., Kincl, L., Krewski, D., McLean, D., Parent, M.-É., Richardson, L., Sadetzki, S., Schlaefer, K., Schlehofer, B., Schüz, J., Siemiatycki, J., van Tongeren, M., & Cardis, E. (2014). Occupational exposure to extremely low-frequency magnetic fields and brain tumor risks in the INTEROCC study. *Cancer Epidemiology, Biomarkers & Prevention, 23*(9), 1863–1872. https://doi.org/10.1158/1055-9965.EPI-14-0102

Vila, J., Turner, M. C., Gracia-Lavedan, E., Figuerola, J., Bowman, J. D., Kincl, L., Richardson, L., Benke, G., Hours, M., Krewski, D., McLean, D., Parent, M.-É., Sadetzki, S., Schlaefer, K., Schlehofer, B., Schüz, J., Siemiatycki, J., van Tongeren, M., & Cardis, E. (2018). Occupational exposure to high-frequency electromagnetic fields and brain tumor risk in the INTEROCC study: An individualized assessment approach. *Environment International, 119*, 353–365. https://doi.org/10.1016/j.envint.2018.06.038

Zhang, F., Xu, C.-L., & Liu, C.-M. (2015). Drug delivery strategies to enhance the permeability of the blood-brain barrier for treatment of glioma. *Drug Design, Development and Therapy*, *9,* 2089–2100. https://doi.org/10.2147/DDDT.S79592

Zhu, Y., Zhu, L., Lan, Y., Sun, C., & Chen, G. (2026). Exposure to hexavalent chromium and 1800 MHz electromagnetic radiation can synergistically induce intracellular DNA damage in mouse embryonic fibroblasts. *Biochemical and Biophysical Research Communications*, *804,* 153360. https://doi.org/10.1016/j.bbrc.2026.153360